

Phytochemical Composition and Antibacterial Activity of *Trachyspermum copticum* L. essential oil, East Azerbaijan, Iran

Haedeh Mobaiyen^{1*}, Mesam Nasarolah Pour¹, Faranak Elmi²

¹Department of Microbiology, Tabriz Branch, Islamic Azad University, Tabriz, Iran;

²Department of Biology, Marand Branch, Islamic Azad University, Marand, Iran.

Received Sep 11, 2016; Accepted Apr 24, 2017

Introduction: The aim of this study was to characterize the chemical composition and antimicrobial activity of *Trachyspermum copticum* essential oil (EO). **Methods:** The chemical composition of seed samples obtained from Mianeh city in East Azerbaijan, was assessed using gas chromatography-mass spectrometry (GC-MS). The antimicrobial activity was evaluated by disc diffusion method against methicillin-resistant *Staphylococcus aureus* (MRSA), other extended-spectrum beta-lactamases (ESBLs) producing, as well as Gram-negative and Gram-positive bacteria. The minimum inhibitory concentration (MIC) value of EO was assessed using agar dilution method. **Results:** Thirteen monoterpene hydrocarbons (57.6%) and oxygenated monoterpenes (42.4%) compounds were identified in the EO, of which, 3 compounds, including thymol, m-cymene, and, γ -terpinene were the major components of the EO with quantities of 41.9, 33.53, and 20.42%, respectively. The EO showed antimicrobial activity against ten microorganisms, especially *Streptococcus sanguis*, *S. aureus* (MRSA strain), and *Klebsiella pneumoniae* (ESBL-producing strain), which was potentially better than tetracycline and kanamycin. **Conclusion:** This study confirmed that EO of *T. copticum* has *in vitro* antimicrobial activity against Gram-negative and Gram-positive bacteria, which has made it an alternative antibacterial agent. J Med Microbiol Infec Dis, 2015, 3 (3-4): 71-74

Keywords: Phytochemicals, *Trachyspermum copticum*, Essential Oil.

INTRODUCTION

Staphylococcus aureus is of great concern in healthcare and community settings, due to involvement in life-threatening infections, and development of resistance to most classes of antimicrobial agents. Methicillin-resistant *Staphylococcus aureus* (MRSA) is a cause of healthcare-associated infections, which had a dramatic increase in number in the 1990s [1], and the recent emergence of MRSA in community-associated infections [2-4] highlights the success of this species as a pathogen and its ability to adapt under pressure from antimicrobial agents.

On the other hand, different reports on extended-spectrum β -lactamases (ESBLs) variants and Metallo-beta-lactamases in the second half of the 1980s and broad geographical distribution of bacteria producing these enzymes, have been considered as another epidemiological phenomenon [5]. Carbapenems are often the last treatment option against ESBL-producing organisms. These organisms have become increasingly resistant to quinolones, aminoglycosides, trimethoprim-sulfamethoxazole, and other antibiotics. Continuous consumption of carbapenems has resulted in the emergence of new classes of Gram-negative bacteria, which is known as superbugs [6].

Multidrug-resistant bacteria have an excessive life-threatening importance, not only in the USA, Europe, and Japan but also in undeveloped countries [7], which have prompted researchers to think of new drugs. Nowadays, due to these problems, new antimicrobial agents and medicinal plants are being investigated as alternatives, especially in

countries with the preferred use of these types of drugs, such as Iran. Essential oils (EOs) of medicinal plants contain very potent natural biologically active agents [8].

Trachyspermum copticum is an annual plant that grows in Ethiopia, Egypt, and India, and is cultivated in Afghanistan, Pakistan, and Iran, which made it as a target of medicinal plants. Geographic distribution of these species in Iran is in Azerbaijan, Isfahan, Fars, Kerman, Sistan and Baluchestan, Khorasan, and Tehran provinces [9]. These provinces have different ecology, which affects the efficiency of plants. Studies by Khosravi *et al.* [10] and Hassanshahian *et al.* [11] showed antifungal and antimicrobial activities of this medicinal plant against clinical isolates, respectively. Thus, the present study was designed to assess antibacterial activity of EO of *T. copticum* (L.) collected from East Azerbaijan province of Iran on Gram-negative and Gram-positive standard strains.

***Correspondence:** Haedeh Mobaiyen

Department of Microbiology, Tabriz Branch, Islamic Azad University, Manzarieh, Soleiman Khater St., Tabriz, Iran, 5157745155.

Email: drhmobaiyen@iaut.ac.ir

Tel: +98 (914) 3005489

Fax: +98 (41) 34781587

MATERIAL AND METHODS

Plant materials. Seeds of *T. copticum* were purchased from markets in Mianeh, East Azerbaijan. The collected materials were air-dried at room temperature (~25°C) in the shade and powdered using a laboratory blender. The studied sample was confirmed as *T. copticum* by a botanist.

Essential oil extraction. One hundred grams of each sample were mixed with 500 ml of distilled water and subjected to hydrodistillation using a Clevenger-type apparatus for 3 h until total recovery of oil. The preparation of the EO was performed three times, and the obtained oils were dried over sodium sulfate, weighed, and stored at 4°C until use.

Gas-chromatography-mass spectrometry (GC-MS). The obtained EOs were analyzed using an Agilent 6890 gas chromatograph-mass spectrometer (GC/MS) fitted with HP-5MS capillary column (30 m×0.25 mm) coupled with an Agilent 5973 mass spectrometer (Agilent Technologies, Palo Alto, Canada). The compounds were identified by matching their recorded mass spectra with the data bank mass spectra (Wiley 7N library).

Strains and growth culture. The antibacterial activity tests included 5 Gram-positive and 5 Gram-negative bacteria acquired from the American Type Culture Collection (ATCC) and the Persian Type Culture Collection (PTCC) including *Enterococcus faecalis* (ATCC 29212), *S. aureus* (ATCC25952), *S. aureus* (ATCC33591) as MRSA, *S. aureus* (ATCC29213), *S. sanguis* (PTCC1449), *Enterobacter aerogenes* (ATCC13048), *Klebsiella pneumoniae* (ATCC700603) as an extended spectrum β -lactamases producing bacteria, *Proteus mirabilis* (ATCC43071), and *Escherichia coli* O157:H7 (purchased from Razi Institute of Iran). These strains were kept at -70°C in Trypticase Soy Broth (TSB) with 20% glycerol, inoculated in Blood Agar (BA), and incubated overnight at 35°C. Subsequently, one colony from each culture was inoculated into TSB and incubated at 35°C for 24 h with shaking (100 rpm) in order to obtain freshly microbial culture suspension (10^8 CFU/ml) for tests.

Agar disc diffusion method. The antimicrobial activity of EOs was determined with disc diffusion method according to Clinical and Laboratory Standards Institute

guidelines [12]. Briefly, bacterial suspensions (10^8 CFU/ml) were spread on Mueller-Hinton Agar (MHA) using sterile cotton swabs. Then, filter paper discs (6 mmØ; Mast, UK) were impregnated with 10 μ l of essential oils of *T. copticum* and were placed on the surface of Petri dishes. Tetracycline (30 μ g/disc) and Kanamycin (30 μ g/disc) (Himedia, India), were used as control. The tests were performed in duplicate for each strain. Antibacterial activity was evaluated by measuring the radius of inhibition zone to the nearest millimeter.

Determination of minimum inhibitory concentration (MIC). The MIC of *T. copticum* EO was determined using agar dilution method [13]. The twofold serial broth dilution of EO was prepared and delivered to series of MHA plates at 45°C, resulted in a final concentration of 512 μ g/ml to 0.5 μ g/ml of essential oils. A standardized suspension of studied bacteria (10^6 CFU) was inoculated into each plate (12 plates for each series), including the plate without EO as a growth control. The plates were incubated at 35°C for 24 h. The microorganisms that were sensitive to EO in the agar plates didn't grow at the inoculation site, where those that were resistant appeared as circular colonies.

RESULTS

Chemical composition of the essential oil. Hydrodistillation of the *T. copticum* seeds yielded EO. Their GC and GC/MS analysis led to identification and quantification of 13 components (Table 1), representing 100% of the total oil. The oil was dominated by monoterpene hydrocarbons (57.63%) and oxygenated monoterpenes (42.37%). The major components of the EO were thymol, m-cymene, and γ -terpinene (41.94, 33.53, and 20.42%), respectively.

Disc diffusion agar and agar dilution methods. In the screening of antibacterial activity of *T. copticum* EO by disc diffusion method, the greatest inhibition zone (50±3 mm) was against *Streptococcus sanguis* PTCC 1449, and the lowest MICs were for *S. aureus* ATCC 25952 and *S. aureus* ATCC 29213 (both 4 μ g/ml, respectively). The results of the antimicrobial assays of EOs of *T. copticum* are summarized in table 2.

Table 1. Chemical composition of essential oil of *T. copticum* collected from East Azerbaijan, Iran

P.N	Name	^a RI	Area%	Case Number
1	α -Thujene	928	0.43	2867-05-2
2	α -Pinene	931	0.4	80-56-8
3	β -Pinene	961	1.47	99-86-5
4	β -Myrcene	979	0.7	123-35-3
5	3-Carene	1005	t	13466-78-9
6	α -Terpinen	1008	0.44	20126-76-5
7	m-Cymene	1013	33.53	535-77-3
8	γ -Terpinen	1047	20.42	99-85-4
9	Terpinolen	1078	0.17	586-62-9
10	4-Terpineol	1161	0.25	20126-76-5
11	Estragole	1177	0.12	140-67-0
12	Sabinol	1179	t	471-16-9
13	Thymol	1266	41.94	89-83-8
	Monoterpene hydrocarbons		57.63	
	Oxygenated monoterpenes		42.37	
	Total		100	

a. Retention indices measured for n-alkanes (c-9 to c-24) on the nonpolar DB-5 column. Tr, traces (<0.1%)

Table 2. Results of MICs and agar disc diffusion tests for tested bacteria against EO of *T. copticum*'s seeds

Test microorganisms	Disk diffusion agar (mm)			MIC (µg/ml) of
	EO	TE	K	EO
<i>Enterococcus faecalis</i> ATCC 29212	33±0.5	26±0.5	25±3	32
<i>Staphylococcus aureus</i> ATCC 25952	45±2	20±1	20±1	4
<i>Streptococcus sanguis</i> PTCC 1449	50±3	30±0.5	16±0.5	16
<i>Staphylococcus aureus</i> ATCC 33591	30±0.5	30±0.5	14±3	8
<i>Staphylococcus aureus</i> ATCC 29213	48±1	20±1	16±1	4
<i>Enterobacter aerogenes</i> ATCC13048	34±0.5	26±0.5	20±1	32
<i>Klebsiella pneumoniae</i> ATCC 700603	34±2	16±1	16±2	128
<i>Escherichia coli</i> ATCC25922	25±2	23±2	17±2	32
<i>Proteus mirabilis</i> ATCC43071	42±1	15±2	22±1	32
<i>Escherichia coli</i> O157:H7	40±3	22±0.5	20±0.5	32

EO: Essential oil; TE: Tetracycline; K: Kanamycin.




Fig. 1. MIC (0.5-512 µg/ml) of EO of *T. copticum*'s seeds by Agar dilution method. 2Z-12Z, one series of diluted EO in agar; control, growth control of diluted microorganisms

According to the results mentioned above, the growth of all tested microorganisms was inhibited by EO of *T. copticum* seeds. Figure 1 shows the results of the tested microorganisms' growth in plates containing serial dilutions of EO in 11 Petri dishes (512-0.05 µg/ml) as well as one plate without EO as a growth control.

DISCUSSION

The amounts of various compounds of EO of *T. copticum* showed differences in samples taken from different part of Iran. In a study by Mahboubi *et al.* (2011), the major components of samples from Kashan city, Iran, were reported to be thymol, γ -terpinene, and o-cymene (45.9, 20.6, and 19%, respectively) [14]. Akbarnia *et al.* (2005) studied 12 samples from different part of Qazvin, Iran, and found that the major components were thymol, γ -terpinene, and p-cymene (41-45, 28-32, and, 16-25%, respectively) [15]. In another study by Haghiroalsadat *et al.* (2011) on *T. copticum* seeds harvested in Yazd province, Iran, it was revealed that thymol (64.9%) and γ -terpinene (11.1%) were the most dominant components of the EO [16]. Rabiey *et al.* (2014), studied on *T. copticum* seeds harvested in Mashad, Iran, and reported that the most dominant components were thymol (57.18%), p-cymene

(22.55%), and γ -terpinene (13.67%) [19]. A glance at the results, shows that the thymol quantity of EO of *T. copticum* obtained from Mashad city [17] was more than those of our study and Yazd, Qazvin, and Kashan [16-18], and γ -terpinene quantity of our study was the same as that of the samples of Mahboubi *et al.*'s study (2011) in Kashan [14].

Based on the results of the antibacterial effect of EO of *T. copticum*, our hypothesis was documented by growth inhibition zone varied from 25 to 50 mm in the disc diffusion method, and MIC varied from 4 to 32 µg/ml against the studied bacteria in agar dilution method. In a study by Shrivatara *et al.* (2012), on antimicrobial potential of Ajwain collected from India, it was shown that inhibition zone for *E. coli* MTCC-443, *Bacillus subtilis* MTCC-441, and *S. aureus* MTCC-3160 were 14.8, 13.6, and 9.9 mm, respectively [18].

Aggarwal *et al.* (2012) studied in Dehradun on 4 species of bacteria obtained from Culture Collection Center, National Culture Laboratory, Pune, India and found that inhibition zone of the oil for *Salmonella typhi*, *E. coli*, *Lactobacillus* and *Bacillus licheniformis* were 40.45, 37.12, 44.54, and 0 mm, respectively [19]. A study by Oroojalian *et al.* (2010) on EO of *Carcum copticum* showed similar MIC results for *E. coli* O157:H7 and *S. aurus* ATCC 25923

(6 and 25 µg/ml, respectively), in comparison with those of our study (32 and 4 µg/ml, respectively) [20].

The EO of *T. copticum* seeds collected from East Azerbaijan showed the same antibacterial activity as the other parts of Iran. Its antimicrobial activity was similar in MRSA strains to other strains of *S. aureus*, which indicated the importance of this medicinal plant as a new therapeutic agent. The antimicrobial activity of this essential oil against ESBLs-producing bacteria was less than MRSA strains and showed that the most activity was against *Streptococcus sanguis* (normal flora of the mouth), which is involved in dental caries.

ACKNOWLEDGEMENT

The authors would like to thank Tabriz Branch, Islamic Azad University for support of this research, and Mr. Amir Talebpour for identification of seeds of the studied medicinal plant.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest associated with this manuscript.

REFERENCES

- Kobayashi N, Alam M, Nishimoto Y, Urasawa S, Uehara N, Watanabe N. Distribution of aminoglycoside resistance genes in recent clinical isolates of *Enterococcus faecalis*, *Enterococcus faecium*, and *Enterococcus avium*. *Epidemiol Infect.* 2001; 126 (2): 197-204.
- Fridkin SK, Hageman JC, Morrison M, Sanza LT, Como-Sabetti K, Jernigan JA, Harriman K, Harrison LH, Lynfield R, Farley MM; Active Bacterial Core Surveillance Program of the Emerging Infections Program Network. Methicillin-resistant *Staphylococcus aureus* disease in three communities. *N Engl J Med.* 2005; 352 (14): 1436-44.
- Hiramatsu K, Okuma K, Ma XX, Hori S, Kapi M. New trends in *Staphylococcus aureus* infections: glycopeptide resistance in hospital and methicillin resistance in the community. *Curr Opin Infect Dis.* 2002; 15 (4): 407-13.
- Wylie JL, Nowicki DL. Molecular epidemiology of community- and health care-associated methicillin-resistant *Staphylococcus aureus* in Manitoba, Canada. *J Clin Microbiol.* 2005; 43 (6): 2830-6.
- Bauernfeind A, Grimm H, Schweighart S. A new plasmidic cefotaximase in a Clinical isolate of *Escherichia coli*. *Infection.* 1999; 18 (5): 294-8.
- Stornburg E, Mack D. Extended-spectrum beta-lactamases implication for clinical microbiology laboratory, therapy and infection control. *J Infect.* 2003; 47 (4): 237-95.
- Gotz F. Staphylococci in colonization and disease: Prospective targets for drugs and vaccines. *Curr Opin Microbiol.* 2004; 7 (5): 477-87.
- Nychas GJE, Tassou CC, Skandamis P. Antimicrobials from herbs and spices. In: Roller SM, editor. *Natural antimicrobials for the minimal processing of foods*. 1st ed. New York: CRC press, Woodhead publishers; 2003; 176-200.
- Fazeli-nasab B, Fooladvand Z. Review on Iranian *Carum copticum* (L.): Composition and Biological Activities. *European J Med Plants.* 2016; 12 (1): 1-8.
- Khosravi AR, Mohammad Salehi R, Yahyaraeyat R, Mokhtari AR, Panahi P. Chemical Composition and Antifungal Activity of *Trachyspermum copticum* Essential Oil Against *Alternaria alternata* (In-Vitro Study). *J Med Plants.* 2015; 1 (53): 82-8.
- Hassanshahian M, Bayat Z, Saeidi S, Shiri Y. Antimicrobial activity of *Trachyspermum ammi* essential oil against human bacterial. *Int J Adv Boil Boiml Res.* 2014; 2 (1): 18-24.
- CLSI. Performance standards for antimicrobial disk susceptibility tests; approved standard-Tenth edition. CLSI document M02-A10. Wayne, PA: Clinical and Laboratory Standards Institute; 2009.
- Winn WC, Allen SD, Janda WM, Koneman E, Procop G, Schreckenberger P, Woods G. Koneman's color atlas and textbook of diagnostic microbiology. 6th ed. Philadelphia: Lippincott, Williams & Wilkins; 2006; 982-3.
- Mahboubi M, Kazempour N. Chemical composition and antimicrobial activity of *Satureja hortensis* and *Trachyspermum copticum* essential oil. *Iran J Microbiol.* 2011; 3 (4): 194-200.
- Akbarinia A, Sefidkon F, Ghalvand A, Tahmasebi Sarvestani Z, Sharifi Asarabdi A. A Study on chemical composition of Ajwain (*Trachyspermum ammi*) essential oil produced in Qazvin. *J Qazvin Univ Med Sci.* 2005; 9 (3): 22-5 [In Persian].
- Haghiroalsadat BF, Vahidi AR., Azimzadeh M, Kalantar SM, Bernard F, Hokmollahi F. Chemical Assessment of Active Ingredients and Antioxidant Effects of *Trachyspermum Copticum's* Seeds harvested in Yazd Province. *J Rafsanjan Univ Med Sci.* 2012; 11 (3): 197-206 [In Persian].
- Rabiey S, Hosseini H, Rezaei M. Use *Carum copticum* essential oil for controlling the *Listeria monocytogenes* growth in fish model system. *Braz J Microbiol.* 2014; 45 (1): 89-96.
- Shrivastava V, Shrivastava G, Sharma R, Mahajan I N, Sharma V, Bhardwaj U. Anti microbial potential of Ajwain (*Trachyspermum copticum*): An Immense Medical Spice. *J Pharm Res.* 2012; 5 (7): 3837-40.
- Aggarwal S, Goyal S. In vitro antimicrobial studies of *Trchyspermum ammi*. *Int J Pharm Bio Sci.* 2012; 3 (4): 64-8.
- Oroojalian F, Kasra-Kermanshahi R, Azizi M, Bassami MR. Synergistic antibacterial activity of the essential oils from three medicinal plants against some important food-borne pathogens by micro dilution method. *Iran J Med Arom Plants.* 2010; 2 (2), 133-46 [In Persian].