

# Prevalence of Typhoidal Salmonella Infections and Associated Risk factors in Kaduna Metropolis, Nigeria

Justina Omotola<sup>1\*</sup>, Innocent Ogbonna<sup>1</sup>, Charles Iheukwumere<sup>2</sup>

<sup>1</sup>Department of Microbiology, College of Science, Federal University of Agriculture, Makurdi, Benue State, Nigeria; <sup>2</sup>Department of Botany, College of Science, Federal University of Agriculture, Makurdi, Benue State, Nigeria

## ARTICLE INFO

## **Original Article**

**Keywords:** Typhoidal *Salmonella*, Enteric fever, Nigeria

Received: 17 Sep. 2020 Received in revised form: 06 Oct. 2020 Accepted: 17 Oct. 2020 **DOI:** 10.29252/JoMMID.8.3.84

\*Correspondence Email: joby4real2link@yahoo.com Tel: +2348101860672 Fax: +2348101860672

## ABSTRACT

Introduction: Salmonella species Salmonella typhi and Salmonella paratyphi, types A, B, and C are the causative agents of enteric fever. This disease continues to pose a severe threat to public health, especially in developing countries. This study investigated the prevalence of typhoidal Salmonella infections and the associated risk factors in Kaduna Metropolis, Nigeria. Methods: A questionnaire was used to obtain information from 250 patients attending four selected government hospitals and 50 healthy individuals as the control. Blood samples from participants were obtained to determine the blood groups and genotypes, and stool specimens were used to isolate typhoidal Salmonella species. Results: Thirty cases were among symptomatic patients and two among controls showing a total prevalence of 10.6%. Among the 32 isolates, 25 were S. typhi (78.1%), and seven were S. paratyphi A (21.9%), revealing an approximate ratio of 4:1. Factors significantly associated with the infection prevalence included age, source of drinking water, and frequency of infection. Blood groups and genotypes were not statistically associated with the infection; however, individuals with blood group O and genotype AA were more commonly infected. Also, females, the age group 20-29, singles, and unemployed, showed more infections. Conclusion: The prevalence, the ratio of typhoidal Salmonella species, and the associated risk factors call for public health and control measures, including the provision of suitable drinking water and improving living and sanitary conditions.

## INTRODUCTION

Enteric fever is a collective term for typhoid and paratyphoid infections caused by Salmonella typhi and Salmonella paratyphi A, B, and C [1]. Salmonella paratyphi A is the most prevalent type, while types B and C are rarely reported [2]. Typhoidal Salmonella bacteria are Gram-negative bacilli transmitted via the fecal-oral route; people acquire infections via ingesting contaminated food or water [1]. These enteric bacteria are restricted to humans [3] and have obtained the ability to overcome mucosal barriers and spread systemically even in immune-competent individuals via the acquisition of virulent factors absent in non-typhoidal Salmonella serotypes [4]. The most important reservoirs of the infection are short-term convalescents or chronic human carriers [5]. If left untreated, enteric fever may result in shock, organ failure, or even death [6].

Evidence gathered so far through genomics and clinical investigations have provided insight into host

restriction and pathogenesis of typhoidal *Salmonella* infections. Research has shown that there might be a possible association between individuals' genetic factors and susceptibility to typhoid and paratyphoid infections [7]. Vaccination, commonly proven efficient in controlling other infectious diseases, is effective against typhoid fever only alongside public health enlightenment [8].

According to the Centre for Disease Control and Prevention (CDC), chlorination of drinking water has dramatically decreased typhoid fever transmission in developed countries like the United States [8]. However, the story is different in developing countries like Nigeria, where people live in poor sanitary conditions and have no access to appropriate health care and suitable drinking water [9]. Genetic components such as blood grouping, Rhesus factor (Rh), and genotypes seem to contribute to the susceptibility and outcome of infectious diseases like typhoid and paratyphoid infections [7, 101.

Enteric fever continues to challenge clinicians, public health scientists, and food safety officials [9]. Therefore, understanding individuals' genetic compositions and associated risk factors will increase our knowledge to prevent and control typhoidal infection in developing countries. This study investigated the prevalence of typhoidal *Salmonella* infections and the associated risk factors in Nigeria.

## MATERIALS AND METHODS

Study Area and Population. This cross-sectional descriptive study was conducted in Kaduna metropolis (Fig. 1), Kaduna State, Nigeria, from July 2018 to February 2019. The study population included 250 patients suspected of enteric fever based on clinical diagnosis and 50 healthy volunteers. The patients from 57 residential areas in Kaduna metropolis (30 in Kaduna North and 27 in Kaduna South) referred to four selected hospitals located in different parts of the metropolis, including General Hospital Sabon Tasha (GHST) representing Kaduna South-east, Gwamna Awan General Hospital (GAGH) Kakuri, Kaduna South-west, Barau Dikko Teaching Hospital (BDTH) Ungwan-Rimi and Yusuf Dantosho Memorial Hospital (YDMH) Tudun Wada. representing Kaduna North-east and west respectively.

**Clinical samples.** The blood samples were collected from all participants in bottles containing ethylenediaminetetraacetic acid (EDTA), and stool samples in clean, dry screw-top containers. The samples were transported to the laboratory (NNPC Industrial Clinic Laboratory, Kaduna).

Ethical approval was obtained from the Ministry of Health and Human Resources, Kaduna State (MOH/ADM/744/VOL.1/483), and the Ethical Committee of Barau Dikko Teaching Hospital Health Research, Kaduna State (Reference Number 18-0034). Written consent was obtained from all participants or their guardians (for the participants  $\leq 15$  years) enrolled in the study.

**Isolation of typhoidal** *Salmonella*. Typhoidal *Salmonella* bacteria were isolated using standard bacteriological procedures. A pinch of stool was enriched in Selenite F broth (HiMedia Laboratories Pvt. Ltd., India) and incubated at 37°C for 6-12 h [14]. A loop full of the enriched broth was cultured by streaking onto the *Salmonella-Shigella* Agar plates (Lab M Limited, Heywood Lancashire, United Kingdom). The plates were incubated aerobically at 37°C for 18-24 h and then examined for characteristic typhoidal *Salmonella* colonies, i.e., smooth, colorless,

or transparent, and flat colonies with a clear halo or black center [14-16]. The identity of the isolates was confirmed as *S. typhi* or *S. paratyphi* A by biochemical tests including Gram staining, Kligler Iron Agar (KIA), and urease test (Titan Biotech Ltd., Rajasthan, India), motility, indole, and lysine decarboxylation or deamination, Methyl Red-Voges Proskauer (MR-VP) test, and citrate utilization test (HiMedia Laboratories Pvt. Ltd., India), catalase and oxidase tests [16-20] (Fig. 2).

**Blood group and genotype.** About 2-5 ml of blood was collected aseptically by vein puncture into EDTA bottles. Blood group and *Rh* factor were determined by mixing a drop of whole blood with anti-A, B, and D blood grouping reagents (Smart Diagnostics Kit, United Kingdom) for 45 s. Agglutination reaction was observed for blood groups and *Rh* reactivity as  $AB^{+/-}$ ,  $A^{+/-}$ ,  $B^{+/-}$  and  $O^{+/-}$ . For genotype determination, gel electrophoresis was employed to separate DNA bands in blood samples using a Uniscope electrophoresis tank for about 15 min. The gels were stained with methylene blue 0.1%, and the migration patterns of DNA were used to identify the genotypes *AA*, *AS*, *SS*, and *AC* against known patterns.

Risk factors for typhoidal Salmonella infections. The investigated risk factors included sociodemographic data (gender, age, marital status, occupational status, type of family and level of education), genetic factors (blood group and genotype), living conditions (the type of toilet and residence, and the number of residents), and other predisposing factors (handwashing practice, traveling, the source of water for general use and drinking, source of food and frequency of infection). A wellstructured questionnaire was used to collect sociodemographic data, clinical information, and predisposing factors.

**Data Analysis.** Data analyses were performed using IBM SPSS version 20 and Chi-square analysis. Results were presented as frequency tables, ratios, percentages, and bar charts. *P*-values less than or equal to 0.05 at 95% confidence level were considered significant for the parameters examined for the associations.

**Clinical samples.** Stool and blood samples were obtained from all participants. Blood samples were collected in bottles containing ethylenediaminetetraacetic acid (EDTA), and stool samples in clean, dry screw-top containers, respectively. The samples were immediately transported to the laboratory (NNPC Industrial Clinic Laboratory, Kaduna).

DOI: 10.29252/JoMMID.8.3.84



Fig 1. Map of Nigeria, Kaduna State, and the study area showing residential regions [11-13]

**Isolation of typhoidal** *Salmonella*. Typhoidal *Salmonella* bacteria were isolated using standard bacteriological procedures. A pinch of stool was enriched in Selenite F broth (HiMedia Laboratories Pvt. Ltd., India) and incubated at 37°C for 6-12 h [14]. Using aseptic laboratory procedures, a loop full of the enriched broth was cultured by streaking onto the *Salmonella Shigella* Agar plates (Lab M Limited, Heywood Lancashire, United Kingdom). The plates were incubated aerobically at 37°C for 18-24 h and then examined for characteristic typhoidal *Salmonella* 

colonies, i.e., smooth, colorless, or transparent, and flat colonies with a clear halo or black center [14-16]. The identity of the isolates was confirmed as either *S. typhi* or *S. paratyphi* A by biochemical tests including Gram staining, Kligler Iron Agar (KIA), and urease test (Titan Biotech Ltd., Rajasthan, India), motility, indole, and lysine decarboxylation or deamination, Methyl Red-Voges Proskauer (MR-VP) test, and citrate utilization test (HiMedia Laboratories Pvt. Ltd., India), catalase and oxidase tests [16-20] (Fig. 2).



## RESULTS

One hundred and sixty-three (163) males and 137 females participated in the study, revealing 15 and 17 cases, respectively (**Table 1**). Table 2 shows the participants' age group distribution and the number of cases in each group. The highest number of participants

(97/300) was among the age group 20-29 years and showed the highest number of isolates (10/32) (**Table 2**). The least number of participants (3/300) were elderly ( $\geq$  60 years), and none was positive.

Participants	Negative Individuals (No.)	S. paratyphi A (%)	S. typhi (%)	Typhoid infections (%)	Number of participants
Male	148	3 (9.4%)	12 (37.5%)	15 (46.9%)	163
Female	120	4 (12.5%)	13 (40.6%)	<b>17</b> (53.1%)	137
Total	268	7 (21.9%)	25 (78.1%)	32 (100%)	300

 Table 2.
 Typhoidal Salmonella infections among different age groups.

Age groups (Year)	Males (No.)	Females (No.)	Negative individuals (no.)	S. paratyphi A (%)	S. typhi (%)	Typhoid infections (%)	Number of Participants
$\leq 9$	27	17	36	2 (6.3%)	6 (18.7%)	8 (25.0%)	44
10-19	19	27	41	1 (3.1%)	4 (12.5%)	5 (15.6%)	46
20-29	50	47	87	2 (6.3%)	8 (25.0%)	<b>10</b> (31.3%)	97
30-39	37	29	62	0	4 (12.5%)	4 (12.5%)	66
40-49	22	13	33	1 (3.1%)	1 (3.1%)	2 (6.2%)	35
50-59	5	4	6	1 (3.1%)	2 (6.3%)	3 (9.4%)	9
$\geq 60$	3	0	3	0	0	0	3
Total	163	137	268	7 (21.9%)	25 (78.1%)	32 (100%)	300

Overall, thirty-two *Salmonella* isolates were obtained in this study, showing a 10.6% prevalence (8.3% for *S. typhi* and 2.3% for *S. paratyphi*) (**Table 3**). Of the 32 *Salmonella* isolates obtained, 25 (78.1%) were *S. typhi*, and 7 (21.9%) were *S. paratyphi* A, showing an approximate ratio of 4:1. Among the 250 patients with a clinical diagnosis, 30 had infections, 24 with *S. typhi*, and 6 with *S. paratyphi* A, while among individuals with no clinical manifestations (controls), two had infections with each of *S. typhi* and *S. paratyphi* A (**Table 3**). North Kaduna and South Kaduna had 11 and 21 cases, respectively (**Table 3**).

Among 32 cases, 22 (68.8%) were among singles (194/300), and 10 (31.2%) belonged to married (106/300).Unemployed participants individuals (162/300) had 20 cases (62.5%), while self-employed (69/300) and employed participants (69/300) had 7 and 5 cases (21.9% and 15.6%), respectively. Participants from nuclear families (252/300) had 25 of the 32 cases, while those from the extended (29/300) and polygamous (19/300) families had 5 and 2 cases, respectively. Eleven cases were from participants with a tertiary level of education (153/300), 11 from individuals with secondary education (65/300), nine from those with no education (51/300), and one from those with primary education (31/300).

Table 4 shows that the most common blood group among 300 participants was  $O^+$  (58.3%). Other blood

J Med Microbiol Infect Dis

groups were B<sup>+</sup> (13.7%), A<sup>+</sup> (9.3%), O<sup>-</sup> (8.3%) and AB<sup>+</sup> (7.7%). Among 32 cases, 21 belonged to individuals with blood group O, followed by the individuals in the blood groups B (8/32), AB (2/32), and A individuals (1/32) (Table 4). The highest infection rate was observed in the *AA* genotype (22/32), followed by *AS* genotype (10/32). Among the rest of the participants, seven had genotype *SS* and one genotype *AC*; none was positive for typhoidal infection (Table 4).

Our survey revealed that most of the infections (30/32) were among participants who used water cisterns (280/300), and the remaining two belonged to pit latrines (20/300). About 53.1% (17/32) of infection belonged to participants who lived in crowded habitats (129/300), and the remaining 15 (46.9%) were among those who lived in more comfortable situations (171/300). Similarly, 11 cases came from residences with  $\leq 5$  inhabitants (135/300), and 21 cases from those with > 5 inhabitants (165/300).

Handwashing practices were rated on a scale of 0 to 5, with 5 as the highest level. No infection was among the 9 participants who were in level 5. The majority of the cases (18/32) belonged to level 2 (156/300), followed by level 1 (35/300) with 6 cases. Four cases belonged to each of level 3 (64/300) and 4 (33/300). Participants who traveled frequently (182/300) had 20 of the 32 cases, followed by those who rarely traveled (61/300), showing

7 cases. Those with no history of travel (57/300) showed the remaining 5 cases.

Participants who used well water for general use (117/300) showed the highest cases (17/32), followed by borehole water users (115/300) that had 9 cases. The remaining 6 cases came from those who used pipe water alone or alongside either borehole or well water.

Similarly, those who drank borehole water mainly alongside alternative sources showed 13 cases, while those who drank sachet water, well, and pipe-borne water as primary sources of drinking water alongside alternatives showed 8, 6, and 3 cases, respectively. The remaining 2 cases belonged to suckling babies who used none of the sources.

Table 3. Distribution of typhoidal Salmonella infections in different hospitals and residential areas

Category of participants	Negative individuals (no.)	S. paratyphi A (no.)	S. typhi (no.)	Total cases	Number of Participants
Patients in Hospitals					
BDTH	44	2	4	6	50
GAGH	80	2	11	13	93
GHST	51	1	5	6	57
YDMH	45	1	4	5	50
Total	220	6	24	30	250
Controls					
Healthy Volunteers	48	1	1	2	50
<b>Residential areas</b>					
Kaduna North	87	3	8	11	98
Kaduna South	181	4	17	21	202
Total	268	7	25	32	300
Percentage (%)	89.3	2.3	8.3	10.6	100

Barau Dikko Teaching Hospital (BDTH), Gwamna Awan General Hospital (GAGH), General Hospital Sabon Tasha (GHST), and Yusuf Dantosho Memorial Hospital (YDMH).

participants (171/300) who consumed Many homemade food alongside foods prepared outside homes showed the highest cases (15/32), followed by those who consumed only homemade food (88/300) with 13 cases. Participants who only consumed foods prepared outside of their homes (35/300) showed 2 cases. The remaining two cases were among the suckling babies (6/300) that consumed neither of the food sources. Clinical information showed that 85.4% of participants (256/300) had previously contracted typhoid fever. Among the remaining 14.6% (44/300) with no history of typhoid fever, four were positive, showing an incidence of 9.1% in this study. About 56.3% of cases (18/32) belonged to participants (160/300) with a history of vaccination against typhoid fever in childhood or adulthood. Also, most of the cases (28/32) were among participants (255/300) who experienced the infection at least once a year.

#### DISCUSSION

The present study investigated the prevalence of typhoidal Salmonella infections and associated risk factors in Kaduna metropolis, Nigeria. The prevalence rate of 10.6% obtained in Kaduna metropolis in this study was lower than 17.2% in Calabar and 24.4% in Owerri, Nigeria, using stool culture and Widal test respectively [21, 22]; 19% in Ethiopia, obtained from both blood culture and Widal test [23], and 62.7% obtained in Karu, Nigeria for S. typhi by Widal test and stool culture [24]. This study's low prevalence rate might be due to the methods used, seasonal variations, or other factors within the state and selected hospitals. The approximate infection 4:1 ratio of S. typhi to S. paratyphi A infection in this study agrees with data in India [2], indicating that S. paratyphi A cases have also increased in Nigeria. Interestingly, in our study, two cases were among healthy volunteers (n=50), showing asymptomatic carriers, suggesting the need to monitor these individuals' disease.

	Negative individuals (no.)	S. paratyphi A (no.)	S. typhi (no.)	Total No. of Isolates (%)	Number of Participants
Blood Group					
A-	4	-	-	-	4 (1.3%)
A+	27	1	-	1 (3.1%)	28 (9.3%)
AB-	4	-	-	-	4 (1.3%)
AB+	21	-	2	2 (6.3%)	23 (7.7%)
B-	1	-	-	-	1 (0.3%)
B+	33	2	6	8 (25.0%)	41 (13.7%)
O-	21	1	2	3 (9.4%)	24 (8.3%)
0+	157	3	15	18 (56.3%)	175 (58.3%)
Genotype					
AA	159	4	18	22 (68.8%)	181 (60.3%)
AS	101	3	7	10 (31.2%)	111 (37.0%)
AC	1	-	-	-	1 (0.3%)
SS	7	-	-	-	7 (2.3%)
Total	268	7	25	32 (100%)	300 (100%)

Table 4. Prevalence of typhoidal Salmonella infections and genetic factors, i.e., blood groups and genotypes

The risk factors found to be statistically associated with the infection were participants' age (p=0.034), source of drinking water (p=0.033), and frequency of infection (p=0.001). Typhoid fever might have a relation with blood phenotypes [25]. However, we found no statistical association between genotypes and blood groups with typhoidal infection in this study. In this study, like in South-east Nigeria, the blood group O comprised the most infected individuals [7].

Females showed a higher infection (53.1%) in this study, like women in Ogume, Owerri, and Karu in Nigeria [7, 22, 24]. Like the results obtained in Owerri, Nigeria, the highest infection rates were in the age group 20-29 (Table 2), showing this age group's vulnerability to enteric fever [22]. Eight of 32 cases (25%) were in the age group 0-9, showing agreement with a high incidence of typhoid fever observed in children  $\leq 15$  years of age [24, 26]. These results suggest that hygiene awareness, especially among those handling children, is necessary. The eating habits of the age group 20-29 might have contributed to the most cases, as there was a significant association between age group and source(s) of food (p=0.001). Contaminated foods and waters and unhygienic food preparation in homes seemed to have most likely exposed these age groups to the infection.

No significant association existed between marital status and typhoidal infection (p=0.054), but singles were more susceptible to infection. The difference might be related to the food source, as singles commonly eat foods prepared outside the home.

In Dakar, Bangladesh, typhoid fever showed a strong relationship with socioeconomic conditions [25]. However, in our study, similar to Ethiopia, no association was found between occupational (p=0.829) or educational status (p=0.149) and typhoidal infection [23], suggesting that occupational status and education

level in developing countries have less effect on exposure to infection than personal hygiene.

In this study, no statistical association was between living conditions and typhoid infection. However, living in group households with poor hygiene showed to be a risk factor for enteric fever [27]. Also, a crowded living condition was strongly associated with typhoid fever [25]. Higher numbers of typhoidal *Salmonella* infections among individuals living in crowded habitats with shared toilets in this study suggested that such conditions may foster the disease. Therefore, improved living conditions, proper sewage disposal, and maintenance of sanitary conditions will no doubt limit the risk of infection.

Unlike a study in Ethiopia, where poor handwashing habit was significantly associated with typhoid infection, no statistical association existed between typhoidal infection and the level of handwashing (p=0.839) in this study [23]. However, no cases in the level 5 handwashing group suggested that proper practice of handwashing could be useful in preventing the disease.

No association existed between the infection and frequency of travel (p=0.991). However, frequent travelers included more cases, suggesting that traveling might increase infection rates via contaminated food and water and poor hygiene conditions.

There was a statistical relationship between drinking water sources and typhoid infections (p=0.033), similar to reports for Dhaka, Bangladesh, Sokoto, Nigeria, and Cameroon [25, 28, 29]. The majority of cases in this study (13/32) were borehole users, followed by sachet water (8/32) and well water (6/32), similar to the study in Cameroon, which showed that borehole or well water as the primary source of drinking water was significantly associated with typhoid fever [29]. In Niger, borehole users had the highest percentage of anti-Salmonella antibodies [30]. Another study in Singida, Tanzania, revealed that land drilled wells as the primary water

source with the proximity to pit latrines very likely resulted in contamination of water [31].

No association existed between vaccination and typhoidal infection (p=0.237). This finding shows that vaccination had no effect on typhoidal infection and corroborate an earlier report that vaccines alone could not eliminate typhoid fever, except when combined with the appropriate public health measures [8,32]. Nevertheless, vaccines should be available, especially in the case of outbreaks [33]. An association between the history of typhoid fever and frequency of infections (p=0.001) in this study revealed the recurring pattern of infection, which also shows that a first-time infection [32].

The prevalence (10.6%), incidence (9.1%), and 4:1 ratio of typhoidal *Salmonella* infection by *Salmonella typhi* and *Salmonella paratyphi* A in this study necessitate intensified efforts for the control and prevention of the infection. The risk factors associated with the infection in the present study also call for public health interventions such as enlightenment of risk factors associated with the disease, provision of portable drinking water, improving living conditions, and ensuring sanitary measures.

## ACKNOWLEDGEMENT

We acknowledge all health facilities and participants involved in the study and The Ministry of Health Kaduna State for their maximum cooperation. We also acknowledge Nigerian National Petroleum Cooperation (NNPC) Industrial Clinic, Kaduna, for their material support.

## **CONFLICT OF INTEREST**

The authors declare there is no conflict of interest associated with this article.

#### REFERENCES

1. cdc.gov [Internet]. Atlanta: Centre for Disease Control and Prevention; c 2019 [cited 2020 Jul 19]. Available from: https://wwwnc.cdc.gov/travel/yellowbook/2020/travel-relatedinfectious-diseases/typhoid-and-paratyphoid-fever.

2. Bharmoria A, Vaish VB, Tahlan AK, Majumder S. Analysis of attributing characteristics of *Salmonella enterica serovar Paratyphi* A, B and C across India, during 6 years (2010 to 2015). J. Med. Microb. Diagn. 2016; 5 (1): 220.

3. Dougan G, Baker S. *Salmonella* enteric serovar Typhi and the pathogenesis of typhoid fever. Annu. Rev. Microbiol. 2014; 68: 317-36.

4. Winter SE, Winter MG, Godinez I, Yang H, Russmann H, Andrew-Polymenis HL, et al. A rapid change in virulence gene expression during the transition from the intestinal lumen into tissue promotes systemic dissemination of *Salmonella*. PLoS Pathog. 2010; 6 (8): 1001060.

5. Wijedoru L, Mallett S, Parry CM. Rapid diagnostic tests for typhoid and paratyphoid (enteric) fever. The Cochrane Database Syst. Rev. 2017 (5): CD008892.

6. Yang J, Barrila J, Roland KL, Ott CM, Nickerson CA. Physiological fluid shear alters the virulence potential of invasive multidrug-resistant non-typhoidal *Salmonella typhimurium* D23580. NPJ Microgravity. 2016; 2: 16021.

7. Otoikhian CSO, Okoror LE. Prevalence of typhoid and paratyphoid, in relation to their genotype among students in Novena University, Ogume. Int. J. Pharm. Med. & Biol. Sci. 2012; 1 (2): 217-24.

8. Date KA, Bentsi-Enchill A, Marks F, Fox K. Typhoid fever vaccination strategies. Vaccine. 2015 Jun 19; 33 Suppl 3: C55–61.

9.Tauxe R, Medalla F, Wise M, Folster J. *Salmonella*, the persistent pathogen. APUA Newsletter. Winter 2014; 32 (3): 5, 12-3.

10. Ali S, Vollard AM, Widjaja S, Surjadi C, Van de Vosse E, Van Dissel JT. PARK2/PACRG polymorphisms and susceptibility to typhoid and paratyphoid fever. Clin. Exp. Immunol. 2006; 144 (3): 425-31.

11. Onwumere M, Dogara MD, Aboh H, Guyuk P, Akutson S, Akuso J. An economic analysis on the viability of harnessing wind energy for power generation in Kaduna State, Nigeria. Sci. J. Energy Eng. 2017; 5 (5): 124-9.

12. Rigasa YA, Badamasi AG, Abdulkarim BI. The evolution of value chains and recycling opportunities in the informal management of municipal solid waste of Kaduna Metropolis, Nigeria. BEST. 2015; 12 (1): 498-505.

13. africaprimenews.com [Internet]. Nigeria: Africa Prime News; c2020 [cited 2020 Oct 3]. Available from: https://www.africaprimenews.com/2018/12/24/development/ni geria-kaduna-awards-5-8billion-contracts-in-kagarko/.

14. Moses IB, Oluduro AO, Isawumi A, Ariyo AB, Fashina CD, Ajayi AO. Antibiotic resistance and molecular characterization of *Salmonella* in diarrhoeal patients' faeces in South-western Nigeria. J. Biol. Agr. Healthcare. 2014; 4 (24): 152-61.

15. Mikoleit ML. Biochemical identification of *Salmonella* and *Shigella* using an abbreviated panel of tests. WHO GFN. 2014; 1-45.

16. microbeonline.com [Internet]. Learn Microbiology Online; c2020 [cited 2020 Aug 15]. Available from: https://microbeonline.com/media-used-culture-identificationsalmonella/.

17. catalog.hardydiagnostics.com [Internet]. Santa Maria: Hardy Diagnostics; c2020 [cited 2020 Aug 15]. Available from:

https://catalog.hardydiagnostics.com/cp\_prod/content/hugo/mil medium.htm#:~:text=MIL%20Medium%20is20used%20to,in %20differentiating%20within%20the%20Enterobacteriaceae.

18. Abdullahi M. Incidence and antimicrobial susceptibility pattern of *Salmonella* species in children attending some hospitals in Kano metropolis, Kano State - Nigeria. BAJOPAS. 2010; 3 (1): 202-6.

19. microbiologyinfo.com [Internet]. Online Microbiology Notes; c2020 [ cited 2020 Aug 15]. Available from:

DOI: 10.29252/JoMMID.8.3.84

#### Typhiod Infections and Associated Risk Factors in Kaduna

https://microbiologyinfo.com/citrate-utilization-test-principlemedia-procedure-and-result/.

20. serc.carleton.edu [Internet]. Montana: Microbial Life Educational Resources; c2016 [cited 2020 Aug 15]. Available from:

https://serc.carleton.edu/microbelife/research\_methods/microsc opy/gramstain.html.

21. Oghenevo OJ, Bassey BE, Yhiler NY, Francis UM, Angela O. Antibiotic resistance in extended spectrum beta-lactamases (Esbls) *Salmonella* species isolated from patients with diarrhoea in Calabar, Nigeria. J. Clin. Infect. Dis. Pract. 2016; 1: 107.

22. Ibegbulam-Njoku PN, Chijioke-Osuji CC, Duru FC. Prevalence of antibody titre in healthy individual and enteric fever patients in Owerri, Nigeria. J. Pub Health Epidemiol. 2014; 6 (6): 192-6.

23. Birhanie M, Tessema B, Ferede G, Endris M, Enawgaw B. Malaria, typhoid fever and their co-infection among febrile patients at a rural health centre in Northwest Ethopia: A cross sectional study. Adv. Med. 2014; 2014: 531074.

24. Abioye JOK, Salome B, Adogo LY. Prevalence of *Salmonella* typhi infection in Karu Local Government Area of Nasarawa State, Nigeria. J. Adv. Microbiol. 2017; 6 (2): 1-8.

25. Banik AK, NurulKabir ATM, Alam J. An association between typhoid fever and age, sex and blood phenotypes ABO and Rh among children – a study in a tertiary care hospital, Dhaka, Bangladesh. IOSR-JDMS. 2018; 17 (9): 63-9.

26. allafrica.com [Internet]. Addis Ababa: Addis Fortune; c2020 [cited 2020 Jul 19]. Available from: https://allafrica.com/stories/201604191347.html.

27. emedicine.medscape.com [Internet]. WebMD LLC; c1994-2020 [cited 2020 Jul 19]. Available from: http://emedicine.medscape.com/article/231135-overview.

28. Ameh IG, Opara WEK. Typhoid: A record of case in Sokoto, Nigeria. Paksitan J. Biol. Sci. 2004; 7 (7): 1177-80.

29. Achonduh-Atijegbe OA, Mfuh KO, Mbange AHE., Chedjou JP, Taylor DW, Nerurkar VR, et al. Prevalence of malaria, typhoid, toxoplasmosis, and rubella among febrile children in Cameroon. BMC Infect. Dis. 2016; 16 (1): 658.

30. Adogo L, Samuel G, Abalaka M. Seroprevalence of *Salmonella* typhi among pregnant women in Niger State. J. Microbiol. Res. 2015; 5 (3): 118-21.

31. Malisa A, Nyaki H. Prevalence and constraints of typhoid fever and its control in an endemic area of Singida region in Tanzania: Lessons for effective control of the disease. J. Pub Health Epidemiol. 2010; 2 (5): 93-9.

32. iamat.org [Internet]. Canada: IAMAT; c2020 [cited 2020 July 19]. Available from: https://www.iamat.org/risks/typhoid-fever.

33. Liston SD, Ovchinnikova OG, Whitfield C. Unique lipid anchor attaches Vi antigen capsule to the surface of *Salmonella* enteric serovar typhi. Proc. Natl. Acad. Sci. USA. 2016; 113 (24): 6719-24.

#### Cite this article:

Omotola J, Ogbonna I, Iheukwumere Ch. Prevalence of Typhoidal *Salmonella* Infections and Associated Risk factors in Kaduna Metropolis, Nigeria. J Med Microbiol Infect Dis, 2020; 8 (3): DOI: 10.29252/JoMMID.8.3.84