

# Journal of Medical Microbiology and Infectious Diseases

JOMM ID ISSN: 2345-5349

eISSN: 2345-5349

# Antioxidant and Antibacterial Activity of *Tanacetum* spp. Essential Oil and Chemical Components

Mostafa Alamholo<sup>1\*</sup>

<sup>1</sup>Department of Biotechnology, Institute of Science and Modern Technology, Rojava University, Qamishlo, Syria

## ARTICLE INFO

# ABSTRACT

## **Original Article**

**Keywords:** *Tanacetum* genus, Infectious bacteria, Phytochemical

Received: 08 May. 2021 Received in revised form: 05 Feb.

2022

Accepted: 16 Feb. 2022 **DOI:** 10.52547/JoMMID.10.1.24

#### \*Correspondence

Email:mostafaalamholo@yahoo.com

Tel: +963938986525

Fax:

© The Author(s)



**Introduction:** The members of the genus *Tanacetum* are important medicinal plants. This study investigated the chemical composition and antibacterial activity of Tanacetum lingulatum and Tanacetum polycephalum essential oils on human infectious bacteria. Methods: The aerial part of two plants were collected from Urmia Province, Iran. The essential oils were extracted using a Clevenger device. The antibacterial effect of essential oils was determined using the disc diffusion assay, minimum inhibitory concentration (MIC), and minimum bactericidal concentration (MBC) by serial dilution method. Also, free radical scavenging activity by 2,2-diphenyl-1-picrylhydrazyl (DPPH) was Chemical composition was measured using the Chromatography-Mass Spectrometry (GCMS). Results: The major constituents in T. polycephalum and T. lingulatum essential oils were 1,8- cineole and camphor, respectively. The highest sensitivity (MIC of 0.312 µg mL<sup>-</sup>) was observed with T. polycephalum against Bacillus subtilis. The lowest IC50 (most potent radical scavenging activity) belonged to T. lingulatum essential oil. Pseudomonas aeruginosa and Streptococcus pyogenes showed resistance to T. lingulatum essential oil. Conclusion: The essential oil of T. polycephalum and T. lingulatum are potential natural antibacterials to treat human pathogenic bacteria and can be used as alternatives to produce antimicrobial agents.

## INTRODUCTION

Some people use herbs with antimicrobial properties for treating infectious diseases [1]. The genus Tanacetum belonging to the Asteraceae family contains 200 species worldwide, of which 26 occur in Iran, comprising medicinal plants [2, 3]. In China, essential oils are used in traditional medicine and traditional ceremonies [4]. Due to bacterial resistance, the high treatment cost of synthetic medications, and the antibiotics side effects, attempts are made to develop new antibacterial agents [5]. Essential oils have a low molecular weight and evaporate rapidly at ambient temperatures. Herbal essential oils have antimicrobial properties, and their applications in pharmacy and medicine have been reported [6]. Essential oils are extracted by several methods, including water and distillation, distillation, steam vacuum chromatography, high-performance liquid and chromatography (HPLC) [6].

Due to the presence of flavonoids, sterols, and terpenoids, the *Tanacetum* species have anti-tumor, antifungal, antibacterial, and anti-migraine activities [7]. Terpenes and parathyroid compounds have anticancer and anti-migraine properties and powerful microbicidal effects against bacteria and fungi [8]. Plants contain high

concentrations of carotenoids, tocopherols, tocotrienols, glutathione, ascorbic acid, and enzymes with antioxidant activity, which protect them from hazardous oxidative damage [9]. Phenolic compounds from medicinal herbs and dietary plants possess various bioactivities and play an essential role in preventing infectious diseases [10, 11]. Secondary metabolites reduce the risk of major chronic diseases (e.g., heart disease, cancer, and diabetes) due to potent antioxidant activities [12]. By inhibiting the polymerization of tubulin and preventing the destructive action of free radicals, flavonoids can play a crucial anticancer role [13]. Also, the anti-*Candida* activity of *T. polycephalum* has been reported [14].

Antiseptic, analgesic, anti-inflammatory, and antimicrobial properties of *T. vulgare* and *T. parthenium* are related to borneol, camphor, and terpenes compounds [15, 16]. Several factors, including harvest time, collection, drying, and storage methods, affect the quantity and quality of essential oils [12].

This study investigated the antibacterial and antioxidant activity of *T. lingulatum* and *T. polycephalum* essential oils on pathogenic bacteria *in vitro* and chemical compositions analysis.

## MATERIALS AND METHODS

The chemical materials, including 2, 2-diphenyl-1-picrylhydrazyl (DPPH), Mueller-Hinton agar (MHA), Nutrient broth (NB), Nutrient agar (NA), and ascorbic acid, were purchased from a commercial company (Merck, Germany), and gentamicin and ciprofloxacin antibiotics were obtained from Padtanteb Company (Iran).

**Preparation of essential oils.** *Tanacetum lingulatum* and *T. polycephalum* were collected from Urmia Province, Iran, in 2015. The essential oil was extracted from aerial parts of collected samples using a Clevenger device at Bu-Ali Sina University (Biotechnology Department Laboratory). Amounts of 100 g of powders were heated in water for 3 h and then was dehydrated by anhydrous sodium sulfate [17].

Bacterial strains. Human pathogenic bacteria including Streptococcus pyogenes (PTCC-1447), Bacillus subtilis (PTCC-1156), Bacillus cereus (PTCC-1247), Micrococcus luteus (ATCC 10987), Enterococcus faecalis (PTCC-1195), Staphylococcus aureus (PTCC-1189), Escherichia coli (ATCC-25922), Shigella boydii (PTCC1744), Salmonella typhi (PTCC-1609), Pseudomonas aeruginosa (PTCC-1181), Enterobacter aerogenes (PTCC-1221), Proteus mirabilis (PTCC-1287), Neisseria meningitides (PTCC-4578), Acinetobacter baumannii (PTCC-4413) and Klebsiella pneumoniae (PTCC-1129) were obtained from the Tehran University, Iran. The bacteria suspension concentration was prepared equal to 0.5 McFarland standard  $(1.5 \times 10^8)$ CFU) [18].

**Disc diffusion assay.** The antibacterial activity of essential oil was performed by disc diffusion assay [6]. Various essential oil concentrations 5, 10, and 20  $\mu$ g mL were prepared in dimethyl sulfoxide (DMSO). A volume of 20  $\mu$ l of the bacterial suspension (1.5 × 10<sup>8</sup> CFU) was cultured on an MHA medium. Then, the 20  $\mu$ l essential oil was spread on discs and incubated at 37 °C for 24 h [19]. Gentamicin (10  $\mu$ g) and ciprofloxacin (5  $\mu$ g) discs were used as positive controls [20] and DMSO as a negative control. The inhibitory zone (mm) formed around each disc was measured, and the data were analyzed by SPSS software.

Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC). The MIC and MBC for *T. lingulatum* and *T. polycephalum* essential oils were determined using the micro-dilution broth method in 96-well plates [21]. Various dilutions of essential oils, i.e., 10, 5, 2.5, 1.25, 0.625, and 0.312 μg mL<sup>-</sup> were prepared. The wells in a 96-well plate were filled with 185 μl of NB medium, and 200 μl of 20 μg mL<sup>-</sup> essential oil was added to the first well plate. Then, 200 μl from the first well plate was added to the second well plate, and the process continued. Finally, 15 μl of bacterial suspension was added to each well plate, followed by incubation at 37 °C for 24 h. The lowest dilution with no

bacteria growth was defined as MIC. To calculate MBC, 5 µl from the plates with no bacterial growth was cultured on an MHA medium and incubated at 37 °C for 24 h. The minimum concentration with no bacterial growth was considered as MBC.

**Free radical scavenging activity.** The free radical scavenging activity was assayed as described by others [22]. Various dilutions of essential oils were prepared, i.e., 2, 4, 6, 8, and  $10~\mu g$  mL. The ascorbic acid and 2, 2- DPPH were used as the standard and reagent. The sample absorption was calculated using a spectrophotometer at 517 nm after 30 min [23]. The IC50 for the essential oil and ascorbic acid were measured. The experiments were tested in triple.

Gas Chromatography-Mass Spectrometry (GCMS). Chemical compositions of essential oils were analyzed by GCMS (Tehran University, Iran). The GCMS analysis was carried out using an Agilent 6890N coupled with an Agilent S973 mass detector, using an HP-5, 30 m column. The instrument was programmed with initial heating at 275 °C for 2 min. The temperature dropped to 120 °C, at an increasing rate of 8 °C /min; then to 285°C, an increasing rate of 3.5°C /min. Injection port temperature was ensured as 350 °C and the helium flow rate at 0.9 ml/min. The samples were injected in split/splitless mode. Solvent delay adjusted for 5 min and 1 μl volume injected [5].

**Statistical analysis.** Data as mean  $\pm$  SD and mean comparison by Duncan test at (P<0.05) in a triple with SPSS software were expressed.

## **RESULTS**

Antibacterial activity. Generally, the sensitivity of tested bacteria against essential oils increased by concentration except for *E. coli*. Moreover, the *T. lingulatum* essential oil did not affect *S. pyogenes, P. aeruginosa*, and *E. faecalis* growth. The *T. polycephalum* essential oil showed a more inhibitory effect than *T. lingulatum* essential oil. Furthermore, the highest sensitivity was exhibited with *T. polycephalum* essential oil (20 µg mL<sup>-</sup>) against *B. subtilis*. Finally, the Grampositive bacteria demonstrated more sensitivity than the Gram-negative bacteria. The antibacterial activity of *T. lingulatum* and *T. lingulatum* essential oil are represented in Table 1.

The MIC and MBC for *T. polycephalum* and *T. lingulatum* essential oil against human pathogenic bacteria are represented in Table 2. The MIC for *T. polycephalum* essential oil on *B. subtillis* was 0.312 μg mL<sup>-</sup>. In addition, the MIC for *T. polycephalum* essential oil on *S. aureus* and *K. pneumoniae* and the MIC for *T. lingulatum* on *B. cereus* was 1.25 μg mL<sup>-</sup>. However, *P. aeruginosa* and *S. pyogenes* exhibited resistance against various dilutions of *T. polycephalum* and *T. lingulatum* essential oils, i.e., 10, 5, 2.5, 1.25, 0.625, and 0.312 μg mL<sup>-</sup>. Furthermore, *T. polycephalum* essential oil showed

more inhibitory activity than *T. lingulatum*. The *T. lingulatum* essential oil had no MIC and MBC with *K. pneumoniae* and *E. faecalis*. The MIC and MBC of

gentamicin on *E. faecalis* was  $0.156 \,\mu g \, mL^{-}$  and MIC and MBC of ciprofloxacin on *E. faecalis* and *E. coli* was  $0.07 \,\mu g \, mL^{-}$ .

Table 1. Antibacterial activity (mm) of T. polycephalum and T. lingulatum essential oils against human pathogenic bacteria

		T. polycephalum (µg mL <sup>-</sup> )			T. lingulatum (μg mL <sup>-</sup> )		Gentamicin (µg mL-)	Ciprofloxacin (µg mL <sup>-</sup> )
Bacteria	20	10	5	20	10	5	10	5
S.boydi	12±0.88 mm	11±0.33 mm	9±0.33 mm	11±0.88 mm	10.5±1.2 mm	8±0.57 mm	29±0.57 mm	29.5±0.33 mm
N.meningitides	14±0.88 mm	12±0.57 mm	-	9±0.22 mm	8±0.66 mm	-	19±0.33 mm	28.5±0.66 mm
A. baumannii	15±0.33 mm	13±0.66 mm	7.5±1.2 mm	10±0.33 mm	9±00 mm	-	20±1 mm	28.5±0.66 mm
K.pneumoniae	17±0.88 mm	$14\pm0.88 \text{ mm}$	9±00 mm	12±0.33 mm	-	-	20±0.57 mm	31.5±0.33 mm
S.typhi	12.5±0.33 mm	11±0.22	8±0.66 mm	15±0.88 mm	$14\pm0.88 \text{ mm}$	12±1.2mm	22±0.33 mm	30±1 mm
P.aeruginosa	8.5±0.33 mm	7±0.66 mm	-	-	-	-	16±0.33 mm	17.5±0.57 mm
E.coli	10.5±0.33 mm	12±0.88 mm	14±0.33 mm	11±0.33 mm	9±0.66 mm	7±0.22 mm	19±0.57 mm	37.5±0.66 mm
E.aerogenes	13±0.88 mm	13±0.88 mm	8±0.66 mm	9±0.33 mm	-	-	20±0.33 mm	24.5±0.66 mm
P. mirabilis	12±0.33 mm	11±0.66 mm	10±0.33 mm	14±0.66 mm	13±0.33 mm	10±0.33mm	19.5±1 mm	24.5±0.57 mm
S.pyogenes	13.5±0.22 mm	12±0.33 mm	10±0.88 mm	-	-	-	11±0.33 mm	28±0.33 mm
M.luteus	18±0.33 mm	15±0.88 mm	13±0.55 mm	15±0.33 mm	12±0.66 mm	10±0.66 mm	15±0.33 mm	17±0.57 mm
S.aureus	18.5±0.33 mm	$16\pm0.66 \text{ mm}$	10±0.33 mm	13±0.88 mm	10±0.33 mm	-	$16\pm0.88 \text{ mm}$	17.5±0.88 mm
B.subtillis	23±0.88 mm	20±0.33 mm	15±0.66 mm	16±0.33 mm	14±00 mm	10±1 mm	15±0.88 mm	16±0.33 mm
B.cereus	20±0.33 mm	19±0.66 mm	16±0.22 mm	14±0.33 mm	10±0.66 mm	9±0.33 mm	15±0.57 mm	16.5±0.57 mm
E. faecalis	16±0.33 mm	12±00 mm	8±0.22 mm	-	-	-	29.5±1 mm	33±0.57 mm

**Table 2.** MIC and MBC of *T. polycephalum* and *T. lingulatum* essential oil against human pathogenic bacteria

Bacteria	T. polycephalum (µg mL.)		T. lingulatum (µg mL <sup>-</sup> )		Gentamicin (µg mL <sup>-</sup> )		Ciprofloxacin (µg mL <sup>-</sup> )	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
S. boydii	5	10	5	10	1.25	2.5	1.25	2.5
N.meningitides	2.5	5	10	10	2.5	5	0.625	1.25
A. baumannii	5	5	5	-	0.625	1.25	0.312	0.625
K. pneumoniae	1.25	5	-	-	0.625	1.25	0.312	0.312
S.typhi	10	-	2.5	5	0.312	0.625	0.156	0.312
P. aeruginosa	-	-	-	-	5	5	2.5	2.5
E. coli	5	10	10	10	2.5	2.5	0.07	0.07
E. aerogenes	2.5	5	10	-	1.25	2.5	0.625	1.25
P. mirabilis	5	5	5	10	5	5	1.25	1.25
S.pyogenes	-	-	-	-	5	5	2.5	2.5
M.luteus	2.5	5	2.5	5	5	5	2.5	5
S. aureus	1.25	10	2.5	5	5	5	2.5	2.5
B. subtillis	0.312	1.25	1.25	5	5	5	2.5	2.5
B. cereus	2.5	5	1.25	5	2.5	5	1.25	2.5
E. faecalis	5	10	-	-	0.156	0.156	0.07	0.07

**Anti-radical activity by DPPH.** The amount of DPPH free radical inhibition increased by essential oil concentrations. A significant difference was observed

between *T. polycephalum* essential oil IC50 and ascorbic acid as the control. The lowest IC50 (most potent radical scavenging activity) was obtained with *T. lingulatum* essential oil (Table 3).

Table3. The DPPH inhibition percentage and IC50 of T. polycephalum and T.lingulatum essential oil

	Inhibition	Inhibition percentage of DPPH in different concentrations (µg mL <sup>-</sup> )				
Organ	2	4	6	8	10	IC50
T. polycephalum	12.53	16.11	19.3	25.74	31.62	$0.94^{a}$
T.lingulatum	24.85	45.31	52.47	70.64	82.88	$0.42^{b}$
Ascorbic acid	25.58	49.02	56.51	79.85	92.04	$0.39^{b}$

The same letters are not significantly different at P < 0.05

**GCMS analysis.** GCMS analysis identified 23 (91%) and 20 (86%) chemicals in *T. polycephalum* and *T. polycephalum*, respectively (Table 4). The main constituents of *T. polycephalum* essential oil were 1,8 cineole (15.7%), camphor (10.94%), alpha-pinene

(5.4%), and trans-sabinene (5.13%), and of *T. lingulatum* essential oil was 1,8 cineole (18.83%), camphor (14.95%), linally propionate (8.5%), and alpha-terpineol (8.02%). The chemical structures of some essential compounds of *Tanacetum* spp. are reflected in Figure 1.

**Table4**. The chemical compositions of *T. polycephalum* and *T.lingulatum* essential oil

T. polycephalum	Compound content (%)	T.lingulatum	Compound content (%)
1.8- Cineole	15.7	1-8.cineole	18.83
1,8- Cineole Camphor	10.94	Camphor	14.95
α –pinene	5.4	Linalyl propionate	8.5
Trans- sabinene	5.136	$\alpha$ –terpineol	8.02
α-Terpineol	4.75	Sabinene	5.81
Berbenol	4.66	Terpinen-4-ol	5.76
Piperitol isomer	4.35	Isobutylbenzene	5.56
2- hexene,4,4,5-trimethyl	3.09	α –pinene	4.17
Chrysanthenyl acetate	3.02	Chrysanthenone	3.87
Borneol	2.37	γ-Eudesmole	3.28
Linalool	2.29	Gamma-terpinene	1.87
Isoborneol	2.13	Eucarvone	1.64
Pinocarvone	2.05	lpha - terpinene	1.48
Trans-caryophyllene	1.96	1-(hydroxymethyl) methylamino adamantine	1.47
Tetrahydropyranyl ether	1.44	Isocaryophillene	1.37
Bornyl acetate	1.68	Endobornyl-acetate	1.34
Beta-pinene	1.67	$\alpha$ -terpineol(p-menth-1-en-8-ol)	1.16
Cyclohept-4-enone	1.62	Globulol	1.03
4-terpineol	1.62	Camphene	1.04
Nerolidol	1.46	Dichloromethane	0.99
Terpinen-4-ol	1.43		
Camphene	1.4		
Sabinene	1.28		




Fig. 1. The chemical structures of important compounds

## DISCUSSION

The chemical compositions, including cineole and camphor with antimicrobial properties, are used to control human pathogenic agents, especially bacteria [24]. The 1,8-cineole, a natural monoterpene, also known as eucalyptol, has anti-inflammatory activity [25].

In this study, the important chemical compositions of T. polycephalum essential oil comprising 1,8 cineole (15.7%), camphor (10.94%), alpha-pinene (5.4%), transsabinene (5.13%),  $\alpha$ -terpineol (4.75%), berbenol (4.66%), piperitol isomer (4.35%), linalool (2.29%), pinocarvone (2.05%), bornyl acetate (1.68%), and camphene (1.4%) were measured. The dominant compounds, included

camphor (18.5%), pinocarvone (31.4%),  $\alpha$ -pinene (9.5%), and bornyl acetate (5.9%), from the aerial part essential oil [26], camphor (59.1%), camphene (14.9%) and 1,8-cineole (10.1%) from flower essential oil [27], linalool (9.72%) from flower essential oil [28] and borneol (28.3%),  $\beta$ -pinene (10.1%),  $\alpha$ -pinene (6.5%), camphene (6%),  $\alpha$ -terpineol (5.16%) and 1,8- cineole (5.1%) [6], from essential oil of *T. polycephalum* were identified. The result of these groups showed a close similarity with the present study. The factors including area, harvest season, climate, and physiological conditions affect types and chemical compositions [29].

Moreover, the chemical compounds including 1,8 cineole (18.83%), camphor (14.95%), linally propionate

(8.5%), alpha-terpineol (8.02%), sabinene (5.81%), and  $\alpha$ -pinene (4.17%) in *T. lingulatum* essential oil were obtained in the present study. The chemical compositions including 1,8- cineole (18.6%) and camphor (13.9%) [28], and  $\alpha$ -pinene (22.876%), 1,8-cineole (21.472%), sabinene (17.902%), 2-pyrrolidinone (7.196%), and camphor (6.794%) [30] have been reported from the aerial part and flower essential oils of *T. lingulatum*, similar to the present study.

The essential oils catalytic action increases permeability in the cell wall and membrane, destroying the pathogens [31]. According to this research, some bacteria, including *S. typhi*, *P. aeruginosa*, and *S. aureus* showed sensitivity against *T. polycephalum* essential oil, especially in high concentrations. There are reports of antibacterial activity of *T. polycephalum* essential oil on *S. typhi*, *S. epidermidi*, *S. saprophiticus*, *P. aeroginosa*, and *S. aureus* [6], similar to the present research.

Based on this research, the amount of DPPH free radical inhibition increased by essential oil concentrations, and the lowest IC50 belonged to *T. lingulatum* essential oil. The free radicals play a pivotal role in some medical conditions, including cancer and cardiovascular diseases, neural disorder, and diabetes [32].

Some Gram-positive bacteria, including *M. luteus, S. aureus*, and *B. subtillis* due to compounds like camphor, sabinene, 1,8 cineole, and linalyl propionate in *T. polycephalum* and *T. lingulatum*, showed more sensitivity to essential oils than tested antibiotics. Finally, antibacterial constituents of *T. polycephalum* and *T. lingulatum* essential oils can be used to develop antimicrobials against pathogenic bacteria.

### **ACKNOWLEDGMENTS**

This study has been extracted from a research project. The corresponding author thanks the Biotechnology Laboratory, Biotechnology Department, Bu-Ali Sina University, Hamadan, Iran.

### CONFLICT OF INTREREST

The authors declare no conflicts of interest associated with this manuscript.

#### REFERENCES

- 1. Wren media, F. Medicinal plants. CTA. 2007; 1-43.
- 2. Goren N, Arda N, Caliskan Z. Chemical characterization and biological activities of the genus tanacetum. J Nat Prod. 2002; 27: 547-658.
- 3. Panday P B A. Textbook of botany angiosperms, taxonomy, anatomy, embryology, baraut. 2004; 303-11.
- 4. Teixeirada Silva JA. Mining the essential oils of the Anthemideae. Afr J Biotechnol. 2004; 3 (12): 706-20.
- 5. Shojaemehr M, Alamholo M, Soltani J. Investigation of antibacterial and antioxidant activity of *Citrus medica* L extract

- on human pathogenic bacteria. Avicenna J Clin Microb Infec. 2020; 7 (1): 8-14.
- 6. Amiri H. Chemical composition, antibacterial and antioxidant activity of the essential oil of *Tanacetum polycephalum* Schutz.Bip. Int J Botany. 2007; 3 (3): 321-4.
- 7. Bagci E, Kocak A. Essential oil composition of two endemic Tanacetum (*T. nitens* Boiss) Grierson and *T. argenteum* (Lam.) Wild. Subsp. argenteum (Asteraceae) taxa, growing wild in Turkey. Ind Crops Prod. 2010; 31: 542-5.
- 8. Hajiakhondi A, Amiri N, Khakighi-sigaroodi F, Rustaiyan A. A new guaianolide from *Tanacetum fruticulosum* ledeb. DARU. 2003; 11 (4): 171-4.
- 9. Yamauchi M, Tsuruma K, Imai S, Nakanishi T, Umigai N, Shimazawa M, et al. Crocetin prevents retinal degeneration induced by oxidative and endoplasmic reticulum stresses via inhibition of caspase activity. Eur J Pharmacol. 2011; 650 (1): 110–19.
- 10. Shih PH, Yeh CT, Yen GC. Anthocyanins induce the activation of phase II enzymes through the antioxidant response element pathway against oxidative stress-induced apoptosis. J Agric Food Chem. 2007; 55: 9427–35.
- 11. Alamhulu M, Nazeri S. The in vitro antibacterial activity of different organs hydroalcoholic extract of *Dendrostellera lesserti*. J of Plant Researches. 2016; 29 (3): 534-42.
- 12. Prior RL, Wu X, Schaich K. Standardized methods for determination of antioxidant capacity and phenolics in foods and dietary supplements. J Agric Food Chem. 2005; 53: 4290–302.
- 13. Lezama R V, Aguilar R T, Ramos R A, Avila E V, Gutierrez MSP. Effect of plantago major on cell proliferation in vitro. J Ethnopharmacol. 2006; 36-42.
- 14. Ghasemi Pirbalouti A, Bahmani M, Avijgan M. Anti-Candida activity of some of the Iranian medicinal plants. Electron J Biotechnol. 2009; 5 (4): 85-8.
- 15. Paulsen E, Christensen LP, Andersen KE. Do monoterpenes released from feverfew (*Tanacetum parthenium*) plants cause airborne Compositae dermatitis. Contact Derm. 2002; 47: 14-18.
- 16. Mazutti M, Mossi AJ, Cansian RL, Corazza ML, Dariva C, Oliveira JV. Chemical profile Boldo (peumus boldus molina) extracts obtained by compressed and antimicrobial activity of carbon dioxide extraction. Braz J Chem Eng. 2008; 25 (34): 427-34
- 17. Kamal GM, Ashraf MY, Hossein A, Shahzadi A, Ghughtai IC. Antioxidant potential of peel essential oil of three Pakistani citrus species: *Citrus reticulate, Citrus sinensis* and *Citrus paradise*. Pak J Bot. 2013; 45 (4): 1449-54.
- 18. Shojaemehr M, Alamholo M. Antibacterial activity of alcoholic and aqueous extracts of various organs of *Citrus medica* on 10 humans pathogenic in vitro. Iran J Med Microbiol. 2019a; 13 (4): 310-20.
- 19. Alamhulu M, Nazeri S. Assessment of the antioxidant and antibacterial effects of stem and leaf alcoholic extracts of *Dendrostellera lesserti*. JMW. 2015a; 7 (4): 289-98.
- 20. Ayoola GA, Johnson OO, Adelowotan T, Aibin IE, Adenipekun E, Odugbemi TO. Evaluation of the chemical constituents and the antimicrobial activity of the volatile oil of *Citrus reticulata* fruit (Tangerine fruit peel) from South West Nigeria. Afr J Biotechnol. 2008; 7 (13): 2223-7.

#### Alamholo

- 21. Alamhulu M, Nazeri S. Investigation antibacterial and antioxidant activities of alcoholic extracts of flower and root *Dendrostellera lesserti* on some human pathogenic bacteria. Avicenna J Clin Med. 2015b; 21 (4): 277-85.
- 22. Stojicevic SS, Stanisiavljevic IT, Velickovic DT, Veljkovic VB, Lazic ML. Comparative screening of the antioxidant and antimicrobial activities of *Sempervivum marmoreum* L. extracts obtained by various extraction techniques. J Serbian Chem. Soc. 2008; 73 (6): 597-600.
- 23. Sahin F, Gulluce M, Daferera D, Sokmen A, Sokmen M, Polissiou M, et al. Biological activities of the essential oils and methanol extract of *Origanum vulgare* ssp. vulgare in the Eastern Anatolia region of Turkey. Food Control. 2004; 15 (7): 549-57.
- 24. Ernestt E, Pittler MH. The efficacy and safety of feverfew (*Tanacetum parthenium*): an update of a systemic review. Public Health Nutr. 2000; 3 (4): 509-14.
- 25. Juergens U R. Anti-inflammatory Properties of the Monoterpene 1.8-cineole: Current Evidence for Co-Medication in Inflammatory Airway Diseases. Drug Res. 2014; 64 (12): 638-46.
- 26. Najafi Gh, Sefidkon F, Mozaffarian V, Zare Maivan H.The essential oil of *Tanacetum polycephalum* schultz-bip. subsp.

- argyrophyllum (K. Koch.) podlech from Iran. J Essent Oil Res. 2007; 1041-2905.
- 27. Nori-Shargh D, Norouzi-Arasi H, Mirza M, Jaimand K, Mohammadi S. Chemical composition of the essential oil of *Tanacetum polycephalum*. Flavour Fragr J. 1999; 14: 105-6.
- 28. Shojaemehr M, Alamholo M. Investigation of human pathogenic bacteria susceptible against essential oil of *Citrus medica* and antioxidant activity *in vitro*. J Herbal Drugs. 2019b; 10 (1): 47-52.
- 29. Afsharypuor S, Jahromy MM. Constituents of the essential oil of *Tanacetum lingulatum* (Boiss.) Bornm. J Essent Oil Res. 2003; 15 (2): 74-6.
- 30. Olamazadeh S, Amjad L, Shahanipour K.Chemical composition and identification of the essential oil of *Tanacetum lingulatum* in Iran. AEB. 2014; 8 (7): 2461-4.
- 31. Mellencamp M. MIC and MBC of oregano essential oil for common livestock pathogens. Ralco Nutrition. 2008; 1-4.
- 32. Devasagayam TPA, Tilak JC, Boloor KK, Sane KS, Ghaskadbi S, Lele RD. Free radicals and antioxidants in human health: current status and future prospects. J Assoc Physicians India. 2004; 52: 794-804.

#### Cite this article: -

Alamholo M. Antioxidant and Antibacterial Activity of *Tanacetum* spp. Essential Oil and Chemical Components. J Med Microbiol Infect Dis, 2022; 10 (1): 24-29. DOI: 10.52547/JoMMID.10.1.24.