NATION PASSEUR

Journal of Medical Microbiology and Infectious Diseases

ISSN: 2345-5349 eISSN: 2345-5330

Molecular Characterizations and Antimicrobial Susceptibility of Extended-Spectrum \(\mathcal{B}\)-lactamase (ESBL) Producing Proteus spp. Clinical Isolates in Babol, Northern Iran

Atena Malaki¹, Elaheh Ferdosi-Shahandashti^{2, 3*}, Amirhosein Maali^{4, 5}, Parisa Sabbagh³, Aynaz Khademian³

¹Department of Biotechnology, Islamic Azad University, Damghan, Iran; ²Department of Medical Biotechnology, School of Medicine, Babol University of Medical Sciences, Babol, Iran; ³Infectious Diseases and Tropical Medicine Research Center, Babol University of Medical Sciences, Babol, Iran; ⁴Department of Immunology, Pasteur Institute of Iran, Tehran, Iran; ⁵Department of Medical Biotechnology, Faculty of Allied Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

ARTICLE INFO

Original Article

Keywords: *Proteus* spp, Betalactamase, Multi-drug resistance, Antibiotic resistance

Received: 10 Dec. 2021 Received in revised form: 21 Jun.

2022

Accepted: 12 Jul. 2022

DOI: 10.52547/JoMMID.10.3.114

*Correspondence

Email: e.ferdosi@mubabol.ac.ir; elaheh.ferdosi@yahoo.com **Tel**: +989113210431

Fax:

© The Author(s)



ABSTRACT

Introduction: Proteus spp. are opportunistic members of Enterobacteriaceae. accounting for 10% of urinary tract infections and other primary clinical infections. They produce extended-spectrum beta-lactamases (ESBL) that can confer resistance to beta-lactam antibiotics. This study aimed to investigate the prevalence, antimicrobial susceptibility, molecular characteristics, and genetic relationship of ESBL-producing *Proteus* spp. clinical isolates in Babol, Northern Iran. Methods: In this cross-sectional study, out of 112 clinical samples, 30 Proteus spp. isolates were identified via specific biochemical assays. According to the Clinical and Laboratory Standards Institute (CLSI) guidelines, antibiotic susceptibility was evaluated using disc diffusion and agar dilution methods, and polymerase chain reaction (PCR) was used to detect bla_{TEM} and bla_{SHV} genes. **Results:** The resistance rate to tetracycline and sulfamethoxazole was highest by disk diffusion and agar dilution. Multiple drug-resistant (MDR) isolates were 86% and 60% in disk diffusion and agar dilution assays. Seven (23.3%) isolates had the bla_{TEM} genes and 18 (60%) bla_{SHV}. Conclusion: ESBL-producing Proteus spp. was highly prevalent, and the blashy was the most common resistance contributing gene. These findings and relatively high resistance to ampicillin demand more care in prescribing antibiotics. Also, the high prevalence of MDR isolates in patients infected with ESBL-producing Proteus spp. requires continuous surveillance.

INTRODUCTION

Proteus spp. are opportunistic members Enterobacteriaceae responsible for 10% of urinary tract infections, cystitis, polio-nephritis, prostatitis, ulcer, eye, and intra-abdominal infections. Proteus vulgaris, Proteus mirabilis, and Proteus penneri are common pathogens affecting immunosuppressed individuals. Also, Proteus members cause ~15% of nephrolithiasis through alkalinization. These bacteria were documented as extended-spectrum beta-lactamase (ESBL)-producers in 1987 [1]. Extended-spectrum β-lactamase (ESBL) producing Enterobacteriaceae is a public health concern worldwide [2, 3]. They produce enzymes responsible for the hydrolysis of oxyimino-beta-lactam antibiotics [4]. The spread of β-lactam antibiotic-resistant isolates occurs by a wide range of ESBL genes, e.g., blaTEM and bla_{SHV} [5, 6]. These narrow-spectrum β -lactamases are located on plasmid cassettes and contribute to resistance to β -lactam antibiotics. The rapid increase in cephalosporin-resistant *Enterobacteriaceae* containing bla_{TEM} and bla_{SHV} genes poses a major therapeutic challenge [6-8].

Misusing antibiotics has led to the spread of multi-drug resistant (MDR) strains, making it a significant challenge for the medical community. The ESBL resistance increases over time. In community-onset, there is a 0.91% yearly increase in ESBL, while in healthcare onsets, it reaches up to 2.31%. In some countries, the phenotypic ESBL production is estimated at 65% in *Enterobacteriaceae* isolates, and in Europe, ESBL antibiotics were the first-line therapy for the associated

Malaki et al.

infectious diseases. Therefore, the first reports of resistance to ESBL antibiotics came from Europe, and it did not take long for similar reports to be received from around the world [9-13]. The gradual evolution of antibiotic-resistant strains has led to the expression of resistance genes in antibiotic-sensitive bacterial strains through gene mutation and horizontal gene transfer, increasing the MDRs prevalence worldwide [14, 15]. The class I integron is a common factor in distributing and spreading antimicrobial resistance. This class carries more than 40 resistance genes related to aminoglycosides, beta-lactams, chloramphenicol, macrolides, sulfonamides, and disinfectants [16].

This study investigated the bla_{TEM} and bla_{SHV} resistance genes in Proteus spp. and their correlations with antibiotic resistance patterns in hospitalized patients.

MATERIAL AND METHODS

The setting, bacterial isolates, and study design. From March 2018 to April 2019, we collected 112 blood samples from the inpatients at Ayatollah Rohani Hospital, Babol, Northern Iran. *Proteus* spp. were identified based on conventional biochemical and microbiological tests, i.e., biotyping assays. All isolates were stored in Luria Bertani broth (Merck, Germany) containing 20% glycerol at -80 °C for further use.

The Ethical Committee of Babol University of Medical Sciences approved this study (Code No.: MUBABOL.REC.1394.162].

Disk diffusion (DD) assay. Susceptibility testing was performed with ten antibiotics by standard disk diffusion (DD) technique according to CLSI (Clinical and Laboratory Standards Institute) standard procedure [17]. The antibiotics included gentamicin (10 μ g), cefepime (30 μ g), amikacin (30 μ g), ciprofloxacin (5 μ g), imipenem (10 μ g), cefotaxime (30 μ g), ampicillin (10 μ g),

piperacillin/tazobactam (30 μ g), sulfamethoxazole (100 μ g), and tetracycline (10 μ g) (MAST Diagnostics, Merseyside, UK). *Escherichia coli* ATCC 25922 was used as positive quality control.

Agar dilution (AD) method. After preparing stock solution from antibiotics according to CLSI 2018 standard [17], 1.5×10^8 CFU/ml of microbial suspensions were cultured on Mueller-Hinton Agar containing the desired antibiotics (MAST Diagnostics, Merseyside, UK) and incubated at 37 °C for 18 to 24 h. A plate containing a medium with no antibiotics was included in assays as the negative control, and results were evaluated according to the CLSI2018 standard table.

Detection of blashy and blatem genes. According to the manufacturer's instructions, DNA extraction from all isolates was performed using a high pure PCR template preparation kit (Roche, Germany). The extracted DNAs were stored at -20 °C for subsequent steps. The ESBLencoding loci, bla_{TEM}, and bla_{SHV} were amplified by conventional PCR using the primers and conditions described by others (Table 1). The 60 µl PCR reactions contained 10 µl of extracted template DNA, 5 µl of 10x buffer, 1.5mM MgCl₂, 0.2mM dNTPs, 50 pMole of each primer (Copenhagen, Denmark), 1.5U of Taq DNA polymerase (Amplicon Co., Denmark) and ddH₂O to the final volume. Amplification was performed in a thermocycler (Corrbet, Australia) (Table 1), and PCR products were electrophoresed in 1.5% agarose gel. The PCR products were sequenced in both directions using the same primers used in amplification in an automated DNA sequencer device (Forster, USA). The standard strain integron-positive Proteus spp. (ATCC1209) was used as a positive control and integron-negative Proteus spp. (ATCC1053) as a negative control. The generated sequences were analyzed at the National Center for Biotechnology Information (NCBI), available at the (http://www.ncbi.nlm.nih.gov/BLAST/) website.

Table 1. Primers and PCR programs for amplifying blatem and blashy genes

PCR Condition					n				
Target	Primer	Primer sequence	Product size (bp)	No. Cycles	Denaturation	Annealing	Extension	Final Extension	Reference
blа _{тем}	TEM-F	5'-ATGAGTATTCAACATTTCCG-3'	851	30	94 °C for 30 S	55 °C for 30 S	72 °C for 1 min	72 °C for 4 min	[32]
	TEM-R	5'-TTAATCAGTGAGGCACCTAT-3'	001						[02]
bla _{SHV}	SHV-F	5'-ATGCGTTATATTCGCCTGTG-3'	735	35	94 °C for 1 min	60 °C for 1 min	72°C for 1 min	72 °C for 10 min	[33]
	SHV-R	5'-TGCTTTGTTATTCGGGCCAA-3'	.35						

RESULTS

Bacterial Isolation. From March 2018 to April 2019, 30 clinical *Proteus* spp. isolates were collected from 30 patients admitted to Ayatollah Rohani Hospital (Babol,

Northern Iran). Other isolates were excluded from the study.

Antibiotic Resistance Profile. MDR was evaluated by DD assay and AD method. All strains were screened for resistance to 10 antimicrobials by DD. The resistance

rates to tetracycline, sulfamethoxazole, ampicillin, cefotaxime, imipenem, gentamicin, cefepime and ciprofloxacin were 90%, 83.3%, 51.7%, 48.2%,17.2%, 13.7%, 10.3% and %3.4, respectively. In contrast, there was no resistance to amikacin and piperacillin/tazobactam. In the DD method, 86.6% of isolates were MDR phenotype. In the agar dilution

method, the resistance rate to sulfamethoxazole, ampicillin, tetracycline, cefotaxime, and cefepime were 80%, 50%, 13.7%, 13.7%, and 10.3%, respectively. No resistance to ciprofloxacin, gentamicin, amikacin, imipenem, and piperacillin/tazobactam was detected (Table 2). Also, in the AD method, 60% of isolates were MDR phenotype (Tables 3 and 4).

Table 2. Frequency of Antibiotic susceptibility pattern of *Proteus* spp. evaluated by disk diffusion (DD) and Agar dilution (AD) methods

Antibiotics	Method	Resistant	Intermediate	susceptible	Total number	P-value
Ciprofloxacin	Disk diffusion	n = 1 (3.4%)	n = 2 (6.8%)	n = 27(90%)	n = 30	NS*
	Agar dilution	n = 0 (0.0%)	n = 1 (3.4%)	n = 29 (96.6%)	n = 30	
Amikacin	Disk diffusion	n = 0 (0.0%)	n = 2 (6.8%)	n = 28 (93.3%)	n = 30	< 0.001
	Agar dilution	n = 0 (0.0%)	n = 0 (0.0%)	n = 30 (100%)	n = 30	
Tetracycline	Disk diffusion	n = 27 (90%)	n = 0 (0.0%)	n = 3 (10.3%)	n = 30	< 0.001
	Agar dilution	n = 4 (13.7%)	n = 23 (76.6%)	n = 3 (10.3%)	n = 30	
Gentamicin	Disk diffusion	n = 4 (13.7%)	n = 0 (0.0%)	n = 26 (86.6%)	n = 30	NS
	Agar dilution	n = 0 (0.0%)	n = 3 (10.3%)	n = 27 (90%)	n = 30	
Cefotaxime	Disk diffusion	n = 14 (48.2%)	n = 9 (31%)	n = 7 (23.3%)	n = 30	NS
	Agar dilution	n = 4 (13.7%)	n = 10 (34.4%)	n = 16 (53.3%)	n = 30	
Ampicillin	Disk diffusion	n = 15 (51.7%)	n = 3 (10.3%)	n = 12 (40%)	n = 30	NS
	Agar dilution	n = 15 (50%)	n = 0 (0.0%)	n = 15 (50%)	n = 30	
Cefepime	Disk diffusion	n = 3 (10.3%)	n = 21 (72.4%)	n = 6 (20%)	n = 30	< 0.05
_	Agar dilution	n = 3 (10.3%)	n = 0 (0.0%)	n = 27 (90%)	n = 30	
Imipenem	Disk diffusion	n = 5(17.2%)	n = 1 (3.4%)	n = 24 (80%)	n = 30	NS
	Agar dilution	n = 0 (0.0%)	n = 5 (17.2%)	n = 25 (83.3%)	n = 30	
Piperacillin tazobactam	Disk diffusion	n = 0 (0.0%)	n = 0 (0.0%)	n = 30 (100%)	n = 30	< 0.001
	Agar dilution	n = 0 (0.0%)	n = 0 (0.0%)	n = 30 (100%)	n = 30	
sulfamethoxazole	Disk diffusion	n = 24 (8.%)	n = 0 (0.0%)	n = 6 (20%)	n = 30	< 0.001
	Agar dilution	n = 24 (80%)	n = 0 (0.0%)	n = 6 (20%)	n = 30	

 Table 3. Multi-drug resistant (MDR) pattern in isolated antibiotic-resistant *Proteus* spp.

Method	MDR	Antibiotics	Resistant sample count (%)	Total number (%)
	Double-resistant	T+ SXT T + CTX + SXT T + CTX T + GM + STX	n = 5 (16.6%) $n = 1 (3.3%)$ $n = 1 (3.3%)$ $n = 4 (13.3%)$	<i>n</i> = 7 (23.3%)
Disk diffusion	Triple-resistant	T + GM + STX T + CTX + SXT T + AP + CTX T + CTX + CPM	n = 4 (13.3%) $n = 1 (3.3%)$ $n = 3 (10%)$ $n = 1 (3.3%)$	n = 9 (30%) (%9.9)
	Quadruple- resistant	T + CTX + AP + SXT $T + CTX + CPM + SXT$	n = 4 (13.3%) n = 2 (6.6%)	$n = 6 (20\%)$ $\begin{array}{c} \infty \\ 97 \\ \square \\ \varepsilon \end{array}$
	Quintuplet- resistant	T + CTX + AP + CP + SXT	n = 3 (10%)	n = 3 (10%)
	Sextuplet- resistant	T + GM + CTX + AP + CP + SXT	n = 1 (3.3%)	n = 1 (3.3%)
	Double-resistant	$\begin{aligned} AP + SXT \\ CTX + CPM \end{aligned}$	n = 10 (33.3%) n = 2 (6.6%)	n = 12 (40%)
Agar dilution		T + AP + SXT	n = 3 (10%)	- 4
r dill	Triple-resistant	CTX + AP + SXT	n = 2 (6.6%)	n = 5 (16.6%)
Aga	Quadruple- resistant	T + CTX + CPM + SXT	n = 1 (3.3%)	n = 5 (16.6%) (% 00 00 00 00 00 00 00 00 00 00 00 00 00


T = Tetracycline; CTX = Cefotaxime; SXT = sulfamethoxazole; GM = Gentamycin; Ap = Ampicillin; CPM = Cefepime; CP = Ciprofloxacin

PCR amplification of Gene Cassettes. Of 30 *Proteus* spp. isolates, 7 (23.3%) and 18 (60%) were positive for bla_{TEM} and bla_{SHV} genes, respectively (Fig. 1).

BLAST and Nucleotide Sequence Accession Number. The positive bla_{TEM} and bla_{SHV} strains were

sequenced and blasted against similar sequences in the Genbank database. After alignment, the homologous sequences were excluded, and the novel ones were deposited in the GenBank database under the accession numbers MH724198, MH724199, MH724200, and MH724201.

a)



b)




Fig. 1. PCR amplification of *bla*_{TEM} (a) and *bla*_{SHV} (b) genes. Lane M: DNA size marker (100bp); C-: negative control (ATCC 1053); C+: positive control (ATCC 1209).

Table 4. Correlation between blaTEM and blaSHV genes and antibiotic resistance

Antibiotics	Antibiotic	D (41	6.0	Ant			
	evolution method	Presence/Abso	ence of Genes	Resistant	Sensitive	Total samples	P-value
	Agar Dilution	TEM	TEM+ TEM-	0 (0%) 1 (4.3%)	7 (100%) 22 (95.7%)	n = 7 n = 23	NS*
ıcin	Dilution	SHV	SHV+	0 (0%)	18 (100%)	n = 23 $n = 18$	NS
Ciprofloxacin	Disk Diffusion	TEM	SHV- TEM+	1 (8.3%) 1 (14.3%)	11 (91.7%) 6 (85.7%)	n = 12 n = 7	NS
			TEM-	2 (8.7%)	21 (91.3%)	n = 23	
J		SHV	SHV+ SHV-	1 (5.6%) 2 (16.7%)	17 (94.4%) 10 (83.3%)	n = 18 n = 12	NS
у <u>к</u> н. В Ъ		TEM	TEM+	0 (0%)	7 (100%)	n = 7	< 0.001

J Med Microbiol Infect Dis 117 2022 Vol. 10 No. 3

						E	SDL in 1 roieus spp.
	Agar		TEM-	0 (0%)	23 (100%)	n = 23	
	Dilution	SHV	SHV+	0 (0%)	18 (100%)	n = 18	< 0.001
		211 /	SHV-	0 (0%)	12 (100%)	n = 12	10.001
	Disk	TEM	TEM+	0 (0%)	7 (100%)	n = 7	NS
	Diffusion		TEM-	2 (8.7%)	21 (91.3%)	n = 23	
		SHV	SHV+	1 (5.6%)	17 (94.4%)	n = 18	NS
			SHV-	1 (8.3%)	11 (91.7%)	n = 12	
	Agar	TEM	TEM+	6 (85.7%)	1 (14.3%)	n = 7	NS
	Dilution	12	TEM-	21 (91.3%)	2 (8.7%)	n = 23	110
(2)		SHV	SHV+	15 (83.3%)	3 (16.7%)	n = 18	NS
Ė			SHV-	12 (100%)	0 (0%)	n = 12	
csc	Disk	TEM	TEM+	6 (85.7%)	1 (14.3%)	n = 7	NS
Tetracycline	Diffusion		TEM-	21 (91.3%)	2 (8.7%)	n = 23	
<u>Te</u>		SHV	SHV+	15 (83.3%)	3 (16.7%)	n = 18	NS
		2111	SHV-	12 (100%)	0 (0%)	n = 12	110
	A	TEM		, ,			NC
	Agar Dilution	TEM	TEM+	0 (0%)	7 (100%)	n = 7	NS
	Dilution		TEM-	3 (13%)	20 (87%)	n = 23	270
3		SHV	SHV+	1 (5.6%)	17 (94.4%)	n = 18	NS
Gentamicin	D: 1	THE A	SHV-	2 (16.7%)	10 (83.3%)	n = 12	NG
31	Disk	TEM	TEM+	0 (0%)	7 (100%)	n=7	NS
3	Diffusion		TEM-	4 (17.4%)	19 (82.6%)	n=23	
		SHV	SHV+	2 (11.1%)	16 (88.9%)	n = 18	NS
			SHV-	2 (16.7%)	10 (83.3%)	n = 12	
	Agar	TEM	TEM+	5 (71.4%)	2 (28.6%)	n = 7	NS
	Dilution		TEM-	9 (39.1%)	14 (60.9%)	n = 23	270
Celotaxillie		SHV	SHV+	10 (55.6%)	8 (44.4%)	n = 18	NS
			SHV-	4 (33.3%)	8 (66.7%)	n = 12	
3	Disk	TEM	TEM+	6 (85.7%)	1 (14.3%)	n = 7	NS
;	Diffusion		TEM-	17 (73.9%)	6 (26.1%)	n = 23	
		SHV	SHV+	15 (83.3%)	3 (16.7%)	n = 18	NS
			SHV-	8 (66.7%)	4 (33.3%)	n = 12	
	Agar	TEM	TEM+	4 (57.1%)	3 (42.9%)	n = 7	NS
	Dilution		TEM-	11 (47.8%)	12 (52.2%)	n = 23	
-		SHV	SHV+	9 (50%)	9 (50%)	n = 18	NS
			SHV-	6 (50%)	6 (50%)	n = 12	
L	Disk	TEM	TEM+	5 (71.4%)	2 (28.6%)	n = 7	NS
Ī	Diffusion	12111	TEM-	13 (56.5%)	10 (43.5%)	n=23	110
		SHV	SHV+	10 (55.6%)	8 (44.4%)	n = 18	NS
		~	SHV-	8 (66.7%)	4 (33.3%)	n = 12	
	Agar	TEM	TEM+	2 (28.6%)	5 (71.4%)	n = 7	NS
	Dilution		TEM-	1 (4.3%)	22 (95.7%)	n = 23	
		SHV	SHV+	3 (16.7%)	15 (83.3%)	n = 18	NS
		~	SHV-	0 (0%)	12 (100%)	n = 12	
Cerebune	Disk	TEM	TEM+	5 (71.4%)	2 (28.6%)	n = 7	NS
3	Diffusion		TEM-	19 (82.6%)	4 (17.4%)	n = 23	
,		SHV	SHV+	14 (77.8%)	4 (22.2%)	n = 18	NS
			SHV-	10 (83.3%)	2 (16.7%)	n = 12	
	Agar	TEM	TEM+	1 (14.3%)	6 (85.7%)	n = 7	NS
	Dilution	I LIVI	TEM-	4 (17.4%)	19 (82.6%)	n=7 $n=23$	145
	Dilation	CITY			` ′		NG
Ę		SHV	SHV+	4 (22.2%)	14 (77.8%)	n = 18	NS
			SHV-	1 (8.3%)	11 (91.7%)	n = 12	
	Disk	TEM	TEM+	1 (14.3%)	6 (85.7%)	n = 7	NS
∃ .	Diffusion		TEM-	5 (21.7%)	18 (78.3%)	n = 23	
		SHV	SHV+	5 (27.8%)	13 (72.2%)	n = 18	NS
			SHV-	1 (8.3%)	11 (91.7%)	n = 12	
Ξ	Agar	TEM	TEM+	0 (0%)	7 (100%)	n = 7	< 0.001
cta.	Dilution		TEM-	0 (0%)	23 (100%)	n = 23	
ğ		SHV	SHV+	0 (0%)	18 (100%)	n = 18	< 0.001
azr			SHV-	0 (0%)	12 (100%)	n = 12	
تہ	Disk	TEM	TEM+	0 (0%)	7 (100%)	n = 7	< 0.001
п	D:cc:		TEM-	0 (0%)	23 (100%)	n = 23	
E E	Diffusion		112111				
Piperacillin tazobactam	Dillusion	SHV	SHV+	0 (0%)	18 (100%)	n = 18	< 0.001

eanı	Agar	TEM	TEM+	5 (71.4%)	2 (28.6%)	n = 7	NS
Co- trim oxa zole	Dilution		TEM-	19 (82.6%)	4 (17.4%)	n = 23	
		SHV	SHV+	14 (77.8%)	4 (22.2%)	n = 18	NS
			SHV-	10 (83.3%)	2 (16.7%)	n = 12	
	Disk	TEM	TEM+	5 (71.4%)	2 (28.6%)	n = 7	NS
	Diffusion		TEM-	19 (82.6%)	4 (17.4%)	n = 23	
		SHV	SHV+	14 (77.8%)	4 (22.2%)	n = 18	NS
			SHV-	10 (83.3%)	2 (16.7%)	n = 12	

DISCUSSION

The prevalence of ESBL- producing Proteus spp. is increasing worldwide, including in the United States, Asia, and Europe [18]. In the present study, the prevalence of bla_{TEM} and bla_{SHV} were 23.3% and 60%. These differences in the distribution of blatem and blashv may be due to geographical distribution, type of organisms, and source of infections. In Iraq, the prevalence of blaTEM was 60% among P. mirabilis isolates [19], while in China, it was around 52% in the same bacteria [20]. In Argentina, investigating the resistance to β-lactam/β-lactamase inhibitors in enterobacteria revealed that all *Proteus* spp. isolates harbored the bla_{TEM} gene [21]. In India, the bla_{TEM} rate among P. mirabilis isolates was 81.9% [22], while in Egypt, it was 35% [23]. In Tehran, Iran, Malekjamshidi et al. (2010) estimated bla_{TEM} prevalence at 83% among ESBL-positive P. mirabilis specimens [24]. Other studies indicated variable rates ranging from 8.3% to 91% [25-29].

In a study by Hamid *et al.*, no *P. mirabilis* isolates in Iraq had *bla*_{SHV} [19]. In India, the *bla*_{SHV} prevalence among *P. mirabilis* isolates was 7% [22]. In Tehran, Iran, the prevalence of *bla*_{SHV} prevalence was 8% in ESBL-positive *P. mirabilis* isolates [24]

The correlation between bla_{TEM} and bla_{SHV} gene and resistance to some antibiotics showed a significant correlation. The bla_{TEM} and bla_{SHV} genes significantly correlated with the resistance to piperacillin/tazobactam obtained by the disk diffusion method. Also, there was a significant correlation between bla_{TEM} and bla_{SHV} genes and resistance to piperacillin/tazobactam and amikacin in the agar dilution method. According to the disk diffusion and agar dilution assays, piperacillin/tazobactam, amikacin, gentamicin, and imipenem are proper choices for treating *Proteus* spp. Given that most ESBL-positive strains showed increased resistance to tetracycline, sulfamethoxazole, cefotaxime, and ampicillin, bla_{TEM} and blashy genes might help confer resistance to these antibiotics. Conza et al. (2014) showed a significant association between the bla_{TEM} gene and resistance to amoxicillin-clavulanic acid [21]. Also, Li et al. (2022) showed a substantial correlation between blaTEM and resistance to chloramphenicol, ciprofloxacin, and trimethoprim-sulfamethoxazole in P. mirabilis isolates [20].

In our study, the highest antibiotic resistance rates were against sulfamethoxazole, tetracycline, and ampicillin. The results of MDR strains in both disk diffusion and agar

dilution methods were 86% and 60%, almost similar to other studies in different countries [17, 26, 27, 30, 31]. Due to the high prevalence of MDR strains, which indicates misuse of antibiotics, studying the physiological properties of β -lactamase genes has received much attention. *Proteus* spp., an opportunistic bacterium, accounts for 10% of urinary tract infections. Therefore, identifying resistance genes is essential for implementing infection control programs and preventing the spread of resistant strains [10-13].

ACKNOWLEDGMENT

The authors thank the Vice Chancellery for Research and Technology of Babol University of Medical Sciences for financial support.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest associated with this manuscript.

REFERENCES

- 1. Malekjamshidi MR, Shahcheraghi F, Feizabadi MM. Detection and PFGE analysis of ESBL-producing isolates of *Proteus* species isolated from patients at Tehran hospitals. Med Sci Monit. 2010; 16 (10): BR327-BR32.
- 2. Peirano G, Pitout JD. Extended-spectrum β-lactamase-producing *Enterobacteriaceae*: update on molecular epidemiology and treatment options. Drugs. 2019: 79 (14): 1529-41.
- 3. Ramadan AA, Abdelaziz NA, Amin MA, Aziz RK. Novel blaCTX-M variants and genotype-phenotype correlations among clinical isolates of extended spectrum beta lactamase-producing *Escherichia coli*. Sci Rep. 2019; 9 (1): 4224.
- 4. Palmeira JD, Ferreira HMN. Extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* in cattle production—a threat around the world. Heliyon. 2020; 6 (1): e03206.
- 5. Cullik A, Pfeifer Y, Prager R, Baum Hv, Witte W. A novel IS26 structure surrounds *blaCTX-M* genes in different plasmids from German clinical *Escherichia coli* isolates. J Med Microbiol. 2010; 59 (5): 580-7.
- 6. Gröbner S, Linke D, Schütz W, Fladerer C, Madlung J, Autenrieth IB, et al. Emergence of carbapenem-non-susceptible extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* isolates at the university hospital of Tübingen, Germany. J Med Microbiol. 2009; 58 (7): 912-22.

- 7. Knudsen PK, Brandtzaeg P, Høiby EA, Bohlin J, Samuelsen Ø, Steinbakk M, et al. Impact of extensive antibiotic treatment on faecal carriage of antibiotic-resistant enterobacteria in children in a low resistance prevalence setting. PLoS One. 2017; 12 (11): e0187618.
- 8. Tumbarello M, Trecarichi EM, Fiori B, Losito AR, D'Inzeo T, Campana L, et al. Multidrug-resistant *Proteus mirabilis* bloodstream infections: risk factors and outcomes. Antimicrob Agents Chemother 2012; 56 (6): 3224-31.
- 9. Vivas R, Barbosa AAT, Dolabela SS, Jain S. Multidrugresistant bacteria and alternative methods to control them: An overview. Microb Drug Resist. 2019; 25 (6): 890-908.
- 10. Drzewiecka D. Significance and roles of *Proteus* spp. bacteria in natural environments. Microb Ecol. 2016; 72 (4): 741-58.
- 11. Armbruster CE, Forsyth-DeOrnellas V, Johnson AO, Smith SN, Zhao L, Wu W, et al. Genome-wide transposon mutagenesis of *Proteus mirabilis*: Essential genes, fitness factors for catheter-associated urinary tract infection, and the impact of polymicrobial infection on fitness requirements. PLoS Pathog. 2017; 13 (6): e1006434.
- 12. Schaffer JN, Pearson MMJ. *Proteus mirabilis* and urinary tract infections. Microbiol Spectr. 2015; 3 (5): 10.
- 13. Chen C-Y, Chen Y-H, Lu P-L, Lin W-R, Chen T-C, Lin C-Y. *Proteus mirabilis* urinary tract infection and bacteremia: risk factors, clinical presentation, and outcomes. J Microbiol Immunol Infect. 2012; 45 (3): 228-36.
- 14. Avlami A, Bekris S, Ganteris G, Kraniotaki E, Malamou-Lada E, Orfanidou M, et al. Detection of metallo-β-lactamase genes in clinical specimens by a commercial multiplex PCR system. J Microbiol Methods. 2010; 83 (2): 185-7.
- 15. Japoni S, Japoni A, Farshad S, Ali AA, Jamalidoust M. Association between existence of integrons and multi-drug resistance in Acinetobacter isolated from patients in southern Iran. Pol J Microbiol. 2011; 60 (2): 163-8.
- 16. Stalder T, Barraud O, Casellas M, Dagot C, Ploy M-C. Integron involvement in environmental spread of antibiotic resistance. Front Microbiol. 2012; 3: 119.
- 17. Chen L, Al Laham N, Chavda KD, Mediavilla JR, Jacobs MR, Bonomo RA, et al. First report of an OXA-48-producing multidrug-resistant *Proteus mirabilis* strain from Gaza, Palestine. Antimicrob Agents Chemother. 2015; 59 (7): 4305-7.
- 18. Uzunović S, Ibrahimagić A, Hodžić D, Bedenić B. Molecular epidemiology and antimicrobial susceptibility of AmpC-and/or extended-spectrum (ESBL) β-lactamase-producing *Proteus* spp. clinical isolates in Zenica-Doboj Canton, Bosnia and Herzegovina. Med Glas (Zenica). 2016; 13 (2): 103-12.
- 19. Fattah Hamid S, Bahadeen Taha A, Jamel Abdulwahid M. Distribution of *blatem*, *blashv*, *blactx-m*, *blaoxa*, and *bladha* in *Proteus mirabilis* Isolated from Diabetic Foot Infections in Erbil, Iraq. Cell Mol Biol (Noisy-le-grand). 2020; 66 (1): 88-94.
- 20. Li Z, Peng C, Zhang G, Shen Y, Zhang Y, Liu C, et al. Prevalence and characteristics of multidrug-resistant *Proteus mirabilis* from broiler farms in Shandong Province, China. Poult Sci. 2022; 101 (4): 101710.

- 21. Di Conza JA, Badaracco A, Ayala J, Rodríguez C, Famiglietti A, Gutkind GO. β-lactamases produced by amoxicillin-clavulanate-resistant enterobacteria isolated in Buenos Aires, Argentina: a new *blatem* gene. Rev Argent Microbiol. 2014; 46 (3): 210-7.
- 22. Chinnam BK, Nelapati S, Tumati SR, Bobbadi S, Chaitanya Peddada V, Bodempudi B. Detection of β-Lactamase-Producing *Proteus mirabilis* Strains of Animal Origin in Andhra Pradesh, India and Their Genetic Diversity. J Food Prot. 2021; 84 (8): 1374-9.
- 23. Algammal AM, Hashem HR, Alfifi KJ, Hetta HF, Sheraba NS, Ramadan H, et al. *atpD* gene sequencing, multi-drug resistance traits, virulence-determinants, and antimicrobial resistance genes of emerging XDR and MDR-*Proteus mirabilis*. Sci Rep. 2021; 11 (1): 9476.
- 24. Malekjamshidi MR, Shahcheraghi F, Feizabadi MM. Detection and PFGE analysis of ESBL-producing isolates of Proteus species isolated from patients at Tehran hospitals. Med Sci Monit. 2010; 16 (10): Br327-32.
- 25. Fouch S, Mitchell J, Lwaleed B, Zinkevich VJ. Evaluation of the shift in antimicrobial resistance due to extended spectrum beta lactamase and AmpC producing *enterobacteriaceae* in Hampshire England. Ann Adv Biomed Sci. 2018; 1 (2): 000109.
- 26. Liakopoulos A, Mevius D, Ceccarelli DJ. A review of SHV extended-spectrum β-lactamases: neglected yet ubiquitous. Front Microbiol. 2016; 7: 1374.
- 27. Fattahi K, Rostamzad AJ. Distribution of *bla_{CTX-M}*, *bla_{TEM}* genes among ESBL producing *Proteus* species isolated from urinary tract infections (UTI) in Ilam. J Res Med Sci. 2015; 39 (1): 41-7.
- 28. Uyanga FZ, Ekundayo EO, Nwankwo EOJ. *bla TEM*, *bla SHV* and *bla CTX-M-15* Extended Spectrum Beta-lactamase Produced by *Acinetobacter baumanii*, *Enterobacter clocae* and *Proteus mirabilis* from Pregnant Women in Three Secondary Health Care Facilities in South-south, Nigeria. J Adv Microbiol. 2019: 1-9.
- 29. Lev AI, Astashkin EI, Kislichkina AA, Solovieva EV, Kombarova TI, Korobova OV, et al. Comparative analysis of *Klebsiella pneumoniae* strains isolated in 2012–2016 that differ by antibiotic resistance genes and virulence genes profiles. Pathog Glob Health. 2018; 112 (3): 142-51.
- 30. Musa HA, Osman MA, Abdelaziz YH, Mohamed S, Ibrahim-Saeed MJ. Distribution of extended-spectrum beta-lactamase *TEM* and *CTX-M* resistance genes among *Proteus* species isolated in Sudan. VacciMonitor. 2019; 28 (2): 80-4.
- 31. Kurihara Y, Hitomi S, Oishi T, Kondo T, Ebihara T, Funayama Y, et al. Characteristics of bacteremia caused by extended-spectrum beta-lactamase-producing *Proteus mirabilis*. J Infect Chemother. 2013; 19 (5): 799-805.
- 32. Schmiedel J, Falgenhauer L, Domann E, Bauerfeind R, Prenger-Berninghoff E, Imirzalioglu C, et al. Multiresistant extended-spectrum β-lactamase-producing *Enterobacteriaceae* from humans, companion animals and horses in central Hesse, Germany. BMC Microbiol. 2014; 14 (1): 187.
- 33. Hujer AM, Page MG, Helfand MS, Yeiser B, Bonomo RA. Development of a sensitive and specific enzyme-linked immunosorbent assay for detecting and quantifying *CMY-2* and *SHV* β-lactamases. J Clin Microbiol. 2002; 40 (6): 1947-57.

Malaki A, Ferdosi-Shahandashti E, Maali A, Sabbagh P, Khademian A. Molecular Characterizations and Antimicrobial Susceptibility of Extended-Spectrum β-lactamase (ESBL) Producing *Proteus* spp. Clinical Isolates in Babol, Northern Iran. J Med Microbiol Infect Dis, 2022; 10 (3): 114-121. DOI: 10.52547/JoMMID.10.3.114