

# Dominant EAEC and Widespread Antibiotic Resistance among Diarrheagenic Escherichia coli in Children with Acute Diarrhea in Zahedan, Iran

Mojdeh Jahantigh<sup>1\*</sup> Nasrin Seyedzayi<sup>2</sup>

<sup>1</sup>Infectious Diseases and Tropical Medicine Research Center, Research Institute of Cellular and Molecular Sciences in Infectious Diseases, Zahedan University of Medical Sciences, Zahedan, Iran; <sup>2</sup>Department of Microbiology, Zahedan University of Medical Sciences, Zahedan, Iran

among children under the age of 10 worldwide, particularly in low- and middle-income countries. Diarrheagenic *Escherichia coli* (DEC) is a

common cause of gastroenteritis in children. This study investigated the

frequency, virulence markers, and antibiotic resistance patterns of DEC in

children below 10 years with acute diarrhea in Zahedan, Iran. Methods: In

this cross-sectional study, 300 *E. coli* isolates were collected from stool samples of children aged below 10 years with diarrhea who presented to hospitals and clinical laboratories in Zahedan. DEC pathotypes were

identified using multiplex PCR and confirmed by standard biochemical tests

and polyvalent antisera. Results: Of the 300 E. coli isolates examined, 89

(29.6%) were identified as diarrheagenic E. coli (DEC) using polyvalent

antisera targeting known DEC pathotypes. Enteroaggregative *E. coli* (EAEC) was identified in 31 isolates (34.83%) based on reaction with

antiserum No. 1. Enterotoxigenic E. coli (ETEC) was identified in 35 isolates

(39.33%) based on reaction with antiserum No. 2. Enteropathogenic E. coli

(EPEC) was identified in 23 isolates (25.84%) based on reaction with antiserum No. 3 (anti-coli3). Multiplex PCR identified the most common pathotype as EAEC (37.6%), followed by EPEC (21.7%), ETEC (15.9%), and EIEC (11.5%). Statistical analysis revealed no significant correlation between the presence of specific virulence genes (*e.g., eae, pcvd432, elt, est,* and *iaH*) and antibiotic resistance patterns in the DEC isolates. **Conclusion**: Given the distribution of DEC pathotypes among children in Zahedan and

their increased antibiotic resistance, antibiotic treatment should be guided by molecular typing and antimicrobial susceptibility testing of isolates, when

# ARTICLE INFO

**Original Article** 

# A B S T R A C T Introduction: Diarrheal disease is a major cause of morbidity and mortality

appropriate.

**Keywords:** Diarrheagenic *E. coli* (DEC), Antibiotic resistance, Enteroaggregative *E. coli* (EAEC), Children, Zahedan, Multiplex PCR

Received: 08 May. 2023 Received in revised form: 30 Mar. 2024 Accepted: 30 Apr. 2024 **DOI:** 10.61186/JoMMID.12.1.59

\*Correspondence Email: mojdehjahantigh001@gmail.com Tel: +989120731775

© The Author(s)

INTRODUCTION

Comprehensive nationwide studies are warranted in Iran to elucidate the distribution of major *E. coli* pathotypes. These studies should prioritize representative sampling strategies and robust study designs to ensure reliable and generalizable findings. Diarrheagenic *E. coli* (DEC) is a common cause of diarrhea in children under five years of age in developing countries, including Iran, where it constitutes a significant public health burden due to its high prevalence and morbidity. In Zahedan, the prevalence of DEC is likely influenced by a combination of factors, including limited access to safe water, inadequate sanitation, and poor hygiene practices, which contribute to the spread of the bacteria [1-4].

DEC strains can be classified into six main pathotypes based on their distinct virulence factors, serotype associations, and epidemiological and clinical characteristics. The pathotypes include enterotoxigenic E. coli (ETEC), enteropathogenic E. coli (EPEC), enteroinvasive E. coli (EIEC), Shiga toxin-producing E. coli (STEC), enteroaggregative E. coli (EAEC), and diffuse adhering E. coli (DAEC), each characterized by specific adhesins, toxins, and other virulence factors that determine their clinical features [5-8].

Each pathotype is defined by a unique combination of virulence factors, which, in conjunction with host factors, influence the clinical manifestations of disease [5]. ETEC is a leading cause of diarrhea in developing countries,

#### Jahantigh et al.

particularly among children under two years old, likely due to factors such as contaminated food and water and limited access to healthcare [6]. In children under one year old, EPEC is a leading cause of diarrhea, accounting for approximately 40-50% of E. coli-related cases. In the period between 2010 and 2020, STEC has been recognized as a significant cause of severe bloody diarrhea, hemorrhagic colitis, and hemolytic uremic syndrome (HUS), particularly among children under five years old. STEC produces a range of virulence factors, including Shiga toxin 1 and Shiga toxin 2, which contribute to its ability to cause severe bloody diarrhea and HUS [9]. EAEC is a significant pathotype associated with a substantial proportion of diarrheal diseases, including endemic and epidemic cases of pediatric diarrhea. EAEC is associated with intestinal inflammation, which can contribute to malnutrition and impaired growth in children, although the relationship is complex and influenced by multiple factors [10, 11]. EAEC and other DEC pathotypes have been identified as important causes of diarrhea in developed countries, contributing to a significant burden of diarrheal diseases [12]. E. coli pathotypes are characterized by their ability to cause severe clinical complications, and their resistance to antibiotics is a significant concern, as it involves a complex interplay of factors, including the type of infectious agent, the location of the microbe, and the concentration of the drug, which are influenced by the patient's immune status [13]. The high prevalence of antibiotic-resistant DEC pathotypes in children, estimated to be around 30-50%, poses a significant public health challenge [14]. The prevalence and pattern of antibiotic resistance for each DEC pathotype are influenced by a complex interplay of factors, including geographic location, health conditions, and socioeconomic status [2].

Accurate identification of diarrheagenic *E. coli* pathotypes and antimicrobial susceptibility testing are essential for guiding targeted and effective treatment strategies, as well as informing evidence-based prevention and control measures [15]. DEC pathotypes are identified using commercial O and H antisera, which detect specific somatic and flagellar antigens, or molecular techniques like M-PCR, which offer higher sensitivity and specificity [16].

Given the significance of diarrheagenic *E. coli* (DEC) in causing diarrhea and the scarcity of comprehensive data in developing countries like Iran, this study aimed to address this knowledge gap. The aim of this study was to determine the prevalence and distribution of DEC pathotypes and their specific antibiotic resistance patterns among children with diarrhea in Zahedan, Iran, to inform evidence-based treatment strategies and identify vulnerable populations.

# MATERIAL AND METHODS

**Sampling and culture.** From July 1, 2016, to October 31, 2016, stool specimens (n=300) were collected from

children with acute diarrhea (defined as three or more loose or watery stools per day) who were either outpatients or inpatients in three hospitals in Zahedan: Ali ibn Abi Talib Hospital, Bu-Ali Hospital, and Khatamalanbeya Hospital. Diarrhea was defined as a history of three or more loose or watery stools per day, or two or more stools of soft or liquid consistency in the 24 h preceding stool collection. Stool samples were processed within 2 hours of collection. The collected stool samples were initially cultured on McConkey Agar, blood Agar, EMB (Merck, Germany), and XLD (Merck, Germany) to isolate E. coli. After 24 h of incubation, the culture plates were examined for the presence of E. coli colonies. Presumptive E. coli colonies were identified using a combination of biochemical tests, including TSI, SIM, MR-VP, Simmons citrate, urea, and lysine decarboxylase. Identification of E. coli isolates was performed using a series of biochemical tests, including fermentation of sugars in TSI medium, indole production and motility in SIM medium, the reaction in MR-VP medium, urea hydrolysis, Simmons citrate utilization, lysine decarboxylation, growth on XLD agar, and growth on EMB medium with a greenish metallic sheen [17].

**Isolation of** *E. coli* **pathotypes by serotyping.** *E. coli* strains were serotyped using Sifin *E. coli* polyvalent antisera and a rapid slide agglutination test. Isolated colonies from 18-h cultures on TSI medium were mixed with one drop of antiserum on a glass slide and incubated for 40 seconds before observation. A positive response was defined as agglutination of bacterial cells upon exposure to the antiserum.

Antibacterial susceptibility testing. Antibiotic susceptibility testing was performed according to CLSI M100-S28 guidelines using the following antibiotics: tetracycline (30 µg), chloramphenicol (30 µg), cotrimoxazole (25 µg), cefixime (5 µg), gentamicin (10 µg), ciprofloxacin (5 µg), imipenem (15 µg), cefotaxime (30 µg), ceftriaxone (30 µg), nalidixic acid (30 µg), ampicillin (10 µg). Antibiotic susceptibility was determined by measuring the diameter of the zone of inhibition around each antibiotic disk, with susceptible isolates showing a zone diameter  $\ge$ X mm, intermediate isolates showing a zone diameter  $\le$ Z mm, according to CLSI criteria [18].

**DNA extraction and multiplex PCR (M-PCR).** Total DNA was extracted from *E. coli* isolates using thermal lysis. The bacterial isolates were cultured for 24 h on LB (Lurria-Bertani) agar. One colony was then transferred to 200  $\mu$ L of sterile water, and the suspension was boiled for 15 min. After the boiling step was complete, the suspension was centrifuged at 13,000 rpm for 10 min. The supernatant was discarded, and the tube was transferred to a laminar flow hood to dry for 30 min. Then, 50  $\mu$ l of distilled water was added to the tube, which was then stored at -20°C. To assess the quality of the purified DNA, 4  $\mu$ l of each DNA sample (dissolved in distilled water)

J Med Microbiol Infect Dis

was electrophoresed on a 1% agarose gel at 80 volts for 45 min. The DNA samples were then stored at -20°C until further use in PCR. The primary PCR mixture was prepared using exact concentrations and volumes of PCR reagents, including MgCl<sub>2</sub> (1.5 mM), dNTPs (200  $\mu$ M), primers (20 pmol), buffer (1X), and distilled water, according to the WHO guidelines for PCR protocols. The PCR mixture was prepared by experienced laboratory personnel trained in molecular biology techniques [18-20].

The PCR reaction was performed in a final volume of 25  $\mu$ L. The reaction mixture were contained 1  $\mu$ L of DNA template, 12  $\mu$ L of Master Mix (Fermentas, Waltham, Massachusetts, United States), 1  $\mu$ M of each primer (Table 1), and 10  $\mu$ L of deionized water (Sigma-Aldrich, USA). PCR assays were performed in a Bio-Rad MJ Mini thermal cycler (T100cycler, Bio-Rad, Hemel Hempstead, UK) with a heated lid for 35 cycles. The PCR products were then analyzed by agarose gel electrophoresis (1% agarose gel stained with ethidium bromide, run for 95 min at 85 volts) for confirmation [18-20].

Target Gene	Primer	Sequence (5 to 3)	Size (bp)	Pathotype
Target Gene eae stx ialH elt	F	TTATGGAACGGCAGAGGT	790	EPEC
	R	CTTCTGCGTACTGCGTTCA		
stx	F	ACGAAATAATTTATATGT	900	STEC
	R	TGATTGTTACAGTCAT		
ialH	F	CTGGTAGGTATGCTGAGG	220	EIEC
	R	CCAGGCCAATTATTTCC	320	EIEC
alt	F	TCTATGTGCATACGGAGC	377	
еп	R	ATACTGATTGCCGCAAT	522	ETEC
est	F	TAAACAAGTAGAGGTCTTCAAAA		
	R	CGGTACAGAGCAGGATTACAACA	147	
D 1/20	F	GGCGAAAGACTGTATCAT	(20)	FAEQ
Pcva432	R ATGTAGAAATCCGCTGTT 630	630	EAEC	

Table 1. Primers used to detect virulence genes in diarrheagenic E. coli by PCR

**Statistical analysis.** Statistical analysis was performed to determine the correlations between the isolated bacterial strains, antibiotic resistance patterns, and specific clinical (symptom severity, disease duration) and demographic variables. The authors performed the statistical analysis using SPSS software version 19. The chi-square test was used to analyze categorical variables, while Fisher's exact test was used to analyze variables with small sample sizes. A *P*-value <0.05 was considered statistically significant.

### RESULTS

**Patient demographics**. Of the 300 samples collected, 173 (57.7%) were from male patients and 127 (42.3%) from female patients. The age of the patients ranged from under nine years and older.

Table 2. Percentage of *E. coli* isolates positive for virulence genes detected using multiplex PCR and confirmed by antiserum agglutination

Identifiable serogroups	Antiserum	Percentage	
O26:K60 O44:K74 O114:K90 O125:K70 O142:k86 O158:K-	Anti-coli1	34.8	
O55:K59 O86:K61 O91:K- O111:K58 O119:K69	Anti-coli2	39.3	
0126:K71 0127:K63 0128:K67 025:K11 078:K80 0103:K- 0118:K- 0124:K72 0145:K- 0157:K- 0164:K-	Anti-coli3	25.8	

Out of the 300 *E. coli* isolates, 89 (29.6%) were identified as diarrheagenic using multiplex PCR and confirmed by antiserum agglutination. Among these 89 isolates, 31 (34.83%) were identified as serogroup O26:K60, O44:K74, O158:K-, O142:K86, O125:K70, and O114:K90 using antiserum No.1 (anti-coli1). Moreover, 35 (29.33%) isolates were identified as

serogroups O55:K59, O128:K67, O127:K63, O126:K71, O119:K69, O111:K58, O91:K-, and O86:K61 using antiserum No. 2 (anti-coli2). Moreover, 23 (25.84%) isolates were identified as serogroups O25:K11, O78:K80, O103:K-, O118:K-, O124:K72, O157:K-, O164:K-, and O145:K- using antiserum No. 3 (anti-coli3).

[ DOI: 10.61186/JoMMID.12.1.59 ]

#### Jahantigh et al.

Antibiotic susceptibility testing. The prevalence of antibiotic resistance among the identified DEC serotypes was evaluated. The highest prevalence of antibiotic resistance was observed for ampicillin (94.8% of all DEC isolates), tetracycline (87.2% of all DEC isolates), and

cotrimoxazole (70.5% of all DEC isolates). The lowest prevalence of antibiotic resistance was observed for imipenem (1% of all DEC isolates) and ciprofloxacin (8.9% of all DEC isolates).



Fig. 1. Antibiotic resistance profile of diarrheagenic *E. coli* (DEC) isolates from children under ten years of age with acute diarrhea in Zahedan, Iran

**Multiplex PCR for pathotypes.** The amplification of the *PCVD432* gene, encoding a virulence factor of Enteroaggregative *E. coli* (EAEC), resulted in a 630-bp amplicon in 26 (37.6%) of the isolated DEC strains. Additionally, the amplification of the *eae* gene, encoding an adhesin of Enteropathogenic *E. coli* (EPEC), yielded an approximately 790-bp amplicon in 15 (21.7%) of the

isolated DEC strains. Among the DEC isolates, 11 (15.9%) carried the *elt* gene, and 9 (13%) carried the *est* gene, both associated with Enterotoxigenic *E. coli* (ETEC). Furthermore, the presence of the *iaH* gene was detected in 8 (11.5%) of the isolates, suggesting their association with Enteroinvasive *E. coli* (EIEC).



Fig. 2. Frequency distribution and relative frequency of ETEC, EPEC, EAEC, and EIEC (%) by age in children under ten years

**Comparing serotyping versus PCR.** Serotyping identified 89 isolates as belonging to specific diarrheagenic *E. coli* pathotypes (EAEC, EPEC, ETEC, EIEC), while molecular analysis using PCR detected 69

isolates carrying specific virulence genes. This discrepancy suggests that serotyping may have a lower detection rate for diarrheagenic *E. coli* compared to PCR.



Fig. 3. Multiplex PCR for identification of EAEC and EPEC pathotypes. Lane1: DNA marker, Lane 2-5: Amplicons specific for EAEC virulence factors, Lane 7-10: Amplicons specific for EPEC virulence factors

### DISCUSSION

Diarrhea is a common childhood illness caused by a variety of enteropathogens, including bacteria, viruses, protozoa. Salmonella, Shigella, Yersinia and enterocolitica, Campylobacter, Vibrio, various E. coli pathotypes, and other bacteria are among the most common causes of bacterial diarrhea [21-23]. E. coli is a significant pathogen that commonly causes watery diarrhea in children under the age of ten, particularly in areas with inadequate access to safe drinking water, such as Zahedan, Iran. Moreover, E. coli has long served as a specific indicator of fecal contamination in water microbiology for over four decades [24]. Estimates suggest that E. coli diarrheal pathotypes are responsible for 30 to 40% of all diarrhea cases, with approximately 20% necessitating hospitalization [17, 25]. Our study included children with acute diarrhea who were referred to hospitals and the central laboratory in Zahedan. Among the 300 E. coli samples analyzed, 89 (29.6%) tested positive for diarrheagenic E. coli (DEC) pathotypes (EAEC, EPEC, ETEC, EIEC) through serotyping and molecular methods (PCR), suggesting a higher prevalence of DEC infection compared to previous studies in the region.

Accurate identification of pathogenic strains is crucial. Therefore, the development and utilization of additional molecular techniques, such as PCR and DNA sequencing, are essential for the reliable detection and characterization of specific diarrheagenic E. coli pathotypes (EAEC, EPEC, ETEC, EIEC) and their virulence factors [26]. As E. coli strains comprise a substantial part of the normal intestinal microflora, distinguishing diarrheagenic E. coli strains from the normal microflora requires the use of phenotypic and genotypic tests that target specific virulence factors and adhesins, such as PCR and DNA sequencing [16]. The present study demonstrated a significantly higher prevalence of diarrheagenic E. coli (29.6%) in children with acute diarrhea in Zahedan, Iran, compared to previous studies in the region [16].

Furthermore, a significant correlation was discovered between the pathotypes and the month of sampling, with the highest frequency observed during the summer months (June to August), particularly in August. This could be attributed to the warmer temperatures (above  $25^{\circ}$ C) and increased humidity, which potentially facilitate increased transmission of bacteria and other pathogenic agents through contaminated water and food, as well as via flies and mosquitoes. This increased vulnerability might be attributed to the developing immune system and the lack of protective antibodies in children under 10 years old. Overall, the causes of diarrhea caused by diarrheagenic *E. coli* in developing countries, especially among children, encompass several factors, including limited access to healthcare facilities, inadequate attention to proper handwashing and food handling practices, absence of proper urban and rural water treatment and sewage systems, and underlying health conditions like malnutrition [14].

The antibiotic resistance patterns in *E. coli* isolates differ among specific communities, countries, and geographical regions, influenced by various factors such as agricultural activities [27]. Tetracycline antibiotics are commonly used to treat human infections, particularly bacterial gastroenteritis, due to their relatively low cost and effectiveness, although they can have significant side effects such as gastrointestinal disturbances and allergic reactions. However, the indiscriminate and inappropriate use of tetracyclines has contributed to the emergence and spread of tetracyclineresistant bacteria, including diarrheagenic *E. coli* [28].

The isolates exhibited relatively low levels of antibiotic resistance against imipenem and ciprofloxacin, respectively. Ciprofloxacin is effective against certain gastrointestinal infections caused by E. coli, such as traveler's diarrhea. Cefixime can be considered as an alternative therapeutic option for the treatment of E. coli-associated diarrhea in pediatric patients when imipenem and ciprofloxacin are ineffective due to resistance. In the present study, the EAEC pathotype was the most prevalent among the diarrheagenic E. coli isolates obtained from children with acute diarrhea, as identified by PCR analysis. Furthermore, PCR analysis identified the presence of the pCVD432 gene, a genetic marker for the EAEC pathotype, in 26 out of 69 isolates (37.6%). Fifteen (21.7%) isolates were identified as EPEC pathotype based on the amplification of the attaching and effacing (eae) gene. The presence of the elt gene and the est gene,

which are genetic markers for the ETEC pathotype, was detected in 11 isolates (15.9%) and 9 isolates (13%), respectively. Furthermore, 8 strains (11.5%) harbored the *iaH* gene, indicating their characterization as the pathotype. Nevertheless, none EIEC of the diarrheagenic E. coli isolates from the children were identified as STEC strains. Significant variations in the frequency of diarrheagenic E. coli pathotypes were observed across different months, with the highest prevalence recorded in August, corresponding to the summer season. A statistically significant association was observed between the prevalence of the four diarrheagenic E. coli pathotypes (EAEC, EPEC, ETEC, and EIEC) and patient age, with the highest proportion occurring in children under 5 years of age, specifically in infants aged 6-8 months. The higher prevalence observed in infants and young children could be attributed to their increased susceptibility to diarrheagenic E. coli infections, likely due to a combination of factors including their developing immune systems, environmental exposure, and host factors. The high burden of diarrheal diseases in developing nations, particularly among children, can be attributed to a complex interplay of factors, including inadequate healthcare infrastructure, poor hand and food hygiene practices, insufficient access to clean water and sanitation, lack of appropriate educational resources, malnutrition, poverty, and environmental factors [27].

In a study conducted by Orni *et al.* (2004), *E. coli* was isolated from 16.9% of the 2,629 children between 1 and 12 years of age with acute diarrhea [29]. Another study carried out in Vietnam in 2002 investigated the prevalence of diarrheal pathogens, which accounted for 27% of all cases examined. Among the identified pathogens, the prevalence of ETEC, EAEC, and EPEC was 6.5%, 15.7%, and 12.2%, respectively [6]. Albert *et al.* (1995) conducted a study on diarrheal and non-diarrheal stool samples collected from children between 1 and 12 years of age in Bangladesh and isolated multiple serotypes, including O125, O127, O114, O126, O55, O128, O142, and O119. Molecular analysis using PCR suggested that most of these serotypes belonged to EPEC strains [30].

In a study conducted in Tabriz, Moghadam *et al.* (2013) identified the most common serotypes using antiserum pool number three, which included O25: K11, O78: K80, O103: K-, O118: K-, O124 K72, O145: K, O157: K-, and O164: K- [16]. In contrast, our study employed antiserum pool number two and identified distinct serotypes.

Haghi *et al.* (2014) reported that children aged 0-5 years had a high prevalence of ETEC pathotype, with a resistance rate of 100% to erythromycin and azithromycin, and a resistance rate of 72.9% to imipenem among children infected with ETEC [31]. In Hamadan, Iran, *E. coli* isolated from children with diarrhea exhibited a high resistance rate to ampicillin

and cotrimoxazole, and a relatively low resistance rate to ciprofloxacin [22]. Diarrheagenic E. coli (DEC) isolates exhibited the highest resistance rates to tetracycline, ampicillin, amoxicillin, amoxicillinclavulanic acid, and trimethoprim [27]. Arif et al. (2010) developed a multiplex PCR assay for diagnosing four major diarrheagenic E. coli pathotypes (ETEC, EPEC, EHEC, and EAEC) by targeting six virulence genes. E. coli was identified in 50 fecal samples using standard biochemical tests such as API 20E or VITEK 2. The multiplex PCR assay detected E. coli target genes in 19 out of 50 (38%) diarrheal samples, indicating the presence of diarrheagenic E. coli. The ETEC group genes (lt or st) were detected in 26.3% (n=5) of the samples, EPEC group genes (eae or bfp) were detected in 63.1% (n=12) of the samples, and EAEC group genes (pCVD) were detected in 21% (n=4) of the samples. Notably, two samples (10.5%) showed co-infection with EAEC group genes, as well as EPEC and ETEC group genes [32]. Lopez et al. (2003) developed a single multiplex PCR assay for identifying gene loci associated with five major diarrheagenic E. coli pathotypes, including EPEC, EAEC, ETEC, and STEC. The PCR assav demonstrated specificity, sensitivity, and rapid detection of target E. coli isolates in fecal and food samples [21].

The prevalence of diarrhea among children admitted to hospitals in both urban and rural hospitals is significantly associated with the presence of diarrheagenic *E. coli* strains. This association was observed in the present study, which is consistent with previous research findings [33-35]. Due to the limited studies on diarrheagenic *E. coli* strains in developing countries, particularly in Iran, it is crucial to perform diagnostic tests in clinical laboratories to further identify these pathogenic strains.

The multiplex PCR assay demonstrates high analytical sensitivity and rapid and accurate detection of specific gene markers associated with diarrheagenic E. coli pathotypes, including EAEC, EPEC, ETEC, and STEC. The simultaneous detection of these pathotypes by the multiplex PCR assay provides a time-saving and cost-efficient alternative that complements conventional phenotypic methods. The multiplex PCR assay can be employed as a valuable tool to complement traditional diagnostic techniques, which are often labor-intensive and cost-prohibitive. Furthermore, due to the increasing antibiotic resistance of these pathotypes, as reported in previous studies, accurate identification and antibiotic susceptibility testing of E. coli strains are essential factors in prescribing effective antimicrobial therapy, in conjunction with other critical considerations such as patient history and clinical presentation. Given the increased antibiotic resistance and mortality rates among hospitalized patients, it is crucial to implement accurate and cost-efficient microbiological diagnostic and therapeutic strategies. These strategies can facilitate

DOI: 10.61186/JoMMID.12.1.59

the identification of diarrheagenic *E. coli* strains and inform the selection of effective antimicrobial therapy.

## ACKNOWLEDGMENT

The authors would like to express their gratitude for the funding provided by Zahedan University of Medical Sciences for this research (Grant ethical code: IR-ZAUMS.REC.1396.156). The data used in this study were de-identified and anonymized to ensure participant privacy and confidentiality. Researchers may obtain access to the data upon legitimate request, with the condition of appropriately citing the present study.

### **CONFLICT OF INTEREST**

The authors declare that there are no conflicts of interest associated with this manuscript.

#### REFERENCES

1. Jafari A, Aslani M, Bouzari S. *Escherichia coli*: a brief review of diarrheagenic pathotypes and their role in diarrheal diseases in Iran. Iran J Microbiol. 2012; 4 (3): 102-17.

2. Schultsz C, Van den Ende J, Cobelens F, Vervoort T, Van Gompel A, Wetsteyn J, et al. Diarrheagenic *Escherichia coli* and acute and persistent diarrhea in returned travelers. J Clin Microbiol. 2000; 38 (10): 3550-4.

3. Bueris V, Sircili MP, Taddei CR, Santos MFd, Franzolin MR, Martinez MB, et al. Detection of diarrheagenic *Escherichia coli* from children with and without diarrhea in Salvador, Bahia, Brazil. Mem Inst Oswaldo Cruz. 2007; 102 (7): 839-44.

4. Omolajaiye S, Afolabi K, Iweriebor B. Pathotyping and antibiotic resistance profiling of *Escherichia coli* isolates from children with acute diarrhea in amatole district municipality of Eastern Cape, South Africa. Biomed Res Int. 2020; 4250165.

5. Podewils LJ, Mintz ED, Nataro JP, Parashar UD. Acute infectious diarrhea among children in developing countries. Semin Pediatr Infect Dis. 2004; 15 (4): 155-68.

6. Hien BTT, Scheutz F, Cam PD, Serichantalergs O, Huong TT, Thu TM, et al. Diarrheagenic *Escherichia coli* and *Shigella* strains isolated from children in a hospital case-control study in Hanoi, Vietnam. J Clin Microbiol. 2008; 46 (3): 996-1004.

7. Zamanzad B, Validi M, Kheiri S, Maghsoudi R. The prevalence and antibacterial susceptibility pattern of Enteropathogenic *Escherichia coli* (EPEC) strains isolated from less than 5 years old children with diarrheal hospitalized in Shahrekord Hajar hospital-2007. J Shahrekord Univ Med Sci. 2010; 11 (4): 11-8.

8. Altwegg M, Perschil I, Gruner E. Molecular biology detection and antibiotic sensitivities of *Shigella* spp. and entero-invasive *Escherichia coli* (EIEC) in patients returning from the tropics. Praxis. 1997; 86 (9): 348-51.

9. Stypułkowska-Misiurewicz H, Augustynowicz E, Andziak J, Kałuzewski S. Susceptibility to vibriostatic factor 0/129 of *Shigella, Salmonella* and entero-invasive *Escherichia coli* (EIEC). Med Dosw Mikrobiol. 1994; 46 (4): 277-83.

10. Gomes TA, Elias WP, Scaletsky IC, Guth BE, Rodrigues JF, Piazza RM, et al. Diarrheagenic *Escherichia coli*. Braz J Microbiol. 2016; 47 Suppl 1(Suppl 1): 3-30.

11. Mansour Amin MS, Javaherizadeh H, Motamedifar M, Saki M, Veisi H, Ebrahimi S, et al. Antibiotic resistance pattern and molecular characterization of extended-spectrum  $\beta$ -lactamase producing enteroaggregative *Escherichia coli* isolates in children from southwest Iran. Infect Drug Resist. 2018; 11: 1097-1104.

12. Vallance B, Finlay B. Exploitation of host cells by enteropathogenic *Escherichia coli*. Proc Natl Acad Sci U S A. 2000; 97 (16): 8799-806.

13. Thapar N, Sanderson IR. Diarrhoea in children: an interface between developing and developed countries. Lancet. 2004; 363 (9409): 641-53.

14. Kumarasinghe G, Lim Y, Chow C, Bassett D. Prevalence of bacterial agents of diarrhoeal disease at the National University Hospital, Singapore and their resistance to antimicrobial agents. Trop Geogr Med. 1992; 44 (3): 229-32.

15. Byrne C, Erol I, Call J, Kaspar C, Buege D, Hiemke C, et al. Characterization of *Escherichia coli* O157: H7 from downer and healthy dairy cattle in the upper Midwest region of the United States. Appl Environ Microbiol. 2003; 69 (8): 4683-8.

16. Zarringhalam M, Nahaei MR, Angouti G, Safaeyan F. Detection of *Escherishia coli* pathotypes and their antibiotic resistance in cases of diarrhea in hospitals of Tabriz in 2013. Res Med. 2015; 38 (4): 233-8.

17. Kaper JB, Nataro JP, Mobley HL. Pathogenic *Escherichia coli*. Nat Rev Microbiol. 2004; 2 (2): 123-40.

18. Zahedani SS, Tahmasebi H, Jahantigh M. Coexistence of Virulence Factors and Efflux Pump Genes in Clinical Isolates of *Pseudomonas aeruginosa*: Analysis of Biofilm-Forming Strains from Iran. Int J Microbiol. 2021; 5557361.

19. Yang J-R, Wu F-T, Tsai J-L, Mu J-J, Lin L-F, Chen K-L, et al. Comparison between O serotyping method and multiplex real-time PCR to identify diarrheagenic *Escherichia coli* in Taiwan. J Clin Microbiol. 2007; 45 (11): 3620-5.

20. Dobrindt U. (Patho-)genomics of *Escherichia coli*. Int J Med Microbiol. 2005; 295 (6-7): 357-71.

21. López-Saucedo C, Cerna JF, Villegas-Sepulveda N, Thompson R, Velazquez FR, Torres J, et al. Single multiplex polymerase chain reaction to detect diverse loci associated with diarrheagenic *Escherichia coli*. Emerg Infect Dis. 2003; 9 (1): 127-31.

22. Alikhani MY, Hashemi SH, Aslani MM, Farajnia S. Prevalence and antibiotic resistance patterns of diarrheagenic *Escherichia coli* isolated from adolescents and adults in Hamedan, Western Iran. Iran J Microbiol. 2013; 5 (1): 42-7.

23. Churgay CA, Aftab Z. Gastroenteritis in children: Part II. Prevention and management. Am Fam Physician. 2012; 85 (11): 1066-70.

24. O'Ryan M, Prado V, Pickering LK. A millennium update on pediatric diarrheal illness in the developing world. Semin Pediatr Infect Dis; 2005: 16 (2): 125-36.

25. Feizizadeh S, Salehi-Abargouei A, Akbari V. Efficacy and safety of *Saccharomyces boulardii* for acute diarrhea. Pediatrics. 2014; 134 (1): e176-e91.

26. Jones ME, Karlowsky JA, Draghi DC, Thornsberry C, Sahm DF, Nathwani D. Epidemiology and antibiotic susceptibility of bacteria causing skin and soft tissue infections in the USA and

J Med Microbiol Infect Dis

Jahantigh et al.

Europe: a guide to appropriate antimicrobial therapy. Int J Antimicrob Agents. 2003; 22 (4): 406-19.

27. Aslani MM, Ahrabi S-A, Alikhani YM, Jafari F, Zali RM, Mani M. Molecular detection and antimicrobial resistance of diarrheagenic *Escherichia coli* strains isolated from diarrheal cases. Saudi Med J. 2008; 29 (3): 388-92.

28. Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. Microbiol Mol Biol Rev. 2001; 65 (2): 232-60.

29. Nahayi M SN, Beygoli N. Genotyping and Antibiotic Susceptibility Patterns of *Pseudomonas aeruginosa* Isolated from Burn Ward of Sina Hospital in Tabriz. Med J Tabriz Uni Med Sciences. 2017; 38 (6): 74-83.

30. Albert MJ, Faruque S, Faruque A, Neogi P, Ansaruzzaman M, Bhuiyan N, et al. Controlled study of *Escherichia coli* diarrheal infections in Bangladeshi children. J Clin Microbiol. 1995; 33 (4): 973-7.

31. Haghi F, Zeighami H, Hajiahmadi F, Khoshvaght H, Bayat M. Frequency and antimicrobial resistance of diarrhoeagenic

*Escherichia coli* from young children in Iran. J Med Microbiol. 2014; 63 (3): 427-32.

32. Arif SK, Salih LI. Identification of different categories of diarrheagenic *Escherichia coli* in stool samples by using multiplex PCR technique. Asian J Me Sci. 2010; 2 (5): 237-43.

33. Shahraki Zahedani S, Jahantigh M, Amini Y. Determining a pattern for antibiotic resistance in clinical isolations of *Pseudomonas aeruginosa*. Tehran Univ Med J. 2018; 76 (8): 517-22.

34. Zelelie TZ, Eguale T, Yitayew B, Abeje D, Alemu A, Seman A, et al. Molecular epidemiology and antimicrobial susceptibility of diarrheagenic *Escherichia coli* isolated from children under age five with and without diarrhea in Central Ethiopia. PloS One. 2023; 18 (7): e0288517.

35. Belete MA, Demlie TB, Chekole WS, Sisay Tessema T. Molecular identification of diarrheagenic *Escherichia coli* pathotypes and their antibiotic resistance patterns among diarrheic children and in contact calves in Bahir Dar city, Northwest Ethiopia. PloS One. 2022; 17 (9): e0275229.

#### Cite this article: -

Jahantigh M, Seyedzayi N. Dominant EAEC and Widespread Antibiotic Resistance among Diarrheagenic *Escherichia coli* in Children with Acute Diarrhea in Zahedan, Iran. J Med Microbiol Infect Dis, 2024; 12 (1): 59-66. DOI: 10.61186/JoMMID.12.1.59.