

Cadmium Bioremoval by *Saccharomyces cerevisiae* in Milk

Ramona Massoud¹, Anoosheh Sharifan^{1*}, Kianoush Khosravi-Darani², GholamHassan Asadi¹

¹Department of Food Science and Technology, Science and Research Branch, Islamic Azad University, Tehran, Iran;

²Department of Food Technology, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Original Article

Keywords: *Saccharomyces cerevisiae*, Cadmium, Bioremoval

Received: Jan. 25, 2020

Received in revised form: Apr. 22, 2020

Accepted: Apr. 27, 2020

DOI: 10.29252/JoMMID.8.1.29

*Correspondence

Email: a_sharifan2000@yahoo.com

Tel: +98 21 44867154

Fax: +98 21 66460700

ABSTRACT

Introduction: The application of biosorbents like bacteria, yeast, and algae is a biotechnological method for eliminating heavy metals from the environment. These microorganisms can also be used for the decontamination of heavy metal in food and water. **Methods:** In this study, we investigated the Cadmium (Cd) biosorption in milk by using *Saccharomyces cerevisiae*. For this purpose, Cd and *S. cerevisiae* were added to milk, and the bioremoval process was monitored for four days. We evaluated six variables, including exposure time, temperature, *S. cerevisiae* concentration, viability yeasts, shaking rate, and initial Cd concentration in the bioremoval process. **Results:** The analysis of ANOVA showed that among the above six variables, *S. cerevisiae* concentration, initial Cd concentration, and exposure time were statistically significantly associated with Cd removal (P values ≤ 0.05). The highest biosorption (70%) was observed after 4 days with 30×10^8 CFU *S. cerevisiae* in milk containing 80 $\mu\text{g/L}$ of Cd. **Conclusion:** Our findings provided further evidence for *S. cerevisiae* as a powerful biosorbent for Cd removal from milk and a potentially safe and green tool for providing safe and healthy food supply.

INTRODUCTION

Heavy metals have adverse effects on human health, posing a threat worldwide [1]. Heavy metals naturally occur in the environment, but anthropogenic activities introduce a large amount of these toxic elements in different environmental compartments resulting in contamination of animals and human food [2]. Worldwide, milk constitutes a central part of the human diet [3, 4]. Different heavy metals may enter the milk supplies through many environmental contaminations like industrial emissions in the air, polluted soil, and contaminated water and plants [5]. Production of heavy metal-free products is one of the primary concerns in the dairy industry, which requires continuous surveillance and adoption of the appropriate method for removing these agents from milk and other dairy product [6]. Cadmium (Cd) is a toxic metal that occurs in industrial wastewaters. The paint, textile, casting, and pharmaceutical industries are significant examples that contribute to heavy metal pollution in the aquatic environment [7]. This heavy metal, even in low concentrations, causes serious health problems to humans, such as hypertension, renal dysfunction, lung damage, teratogenic effects, and hepatic injury [8-10]. Some countries such as Iraq [11], Turkey [12], Nigeria [13], China [14], and Iran [15, 16] have reported Cd contamination in milk.

Various chemical and physical approaches, including precipitation, ultrafiltration, electrolysis, ion exchange, and

reverse osmosis, are available for the removal of heavy metals from aqueous solutions [17]. However, in recent years alternative strategies like bioremediation methods have attracted increasing attention. Some studies have indicated using bacteria, yeasts, and fungi such as *Lactobacillus plantarum*, *Lactobacillus* spp., and *Saccharomyces cerevisiae* for bioremoval of heavy metals like Cd and lead (Pb) from foodstuffs. This novel method is cheaper and more suitable for food and water decontamination [18-20].

In the food industry, *S. cerevisiae* is a familiar and promising candidate for heavy metals biosorption from drinks and food [21, 22]. Some studies have reported the bioremoval of heavy metals by this organism [23-26]. This valuable biosorbent is an available, safe, cheap, and suitable agent for the decontamination of foodstuff [27]. Bioremoval is an absorption process in which heavy metals are attached to the cell surface [28]. In general, heavy metals bioremoval occurs through different mechanisms. The functional groups (hydroxyl and carboxyl) in the cell wall of *S. cerevisiae* play an essential role in the bioremoval process. They are responsible for metal ions fixation during the process. Also, the intracellular metal accumulation happens in the cell membrane, and metal ions can bind to other cellular molecules [27].

There is no much data on Cd bioremoval from milk. This

study, for the first time, reports the use of *S. cerevisiae* for removing low levels ($\mu\text{g/L}$) of Cd concentration from milk. We also investigated *S. cerevisiae* concentration, temperature, exposure time, and heavy metal concentration as the most crucial variables in the bioremoval process [28].

MATERIAL AND METHODS

The Cd removal was performed by various concentrations of *S. cerevisiae* (10 , 20 , 30 , and 40×10^8 CFU/mL) and Cd (20 , 40 , 60 , 80 , 100 mg/L), and exposure times ranging from 1 to 4 days.

Yeast growth. The freeze-dried *S. cerevisiae* PTCC 5020 was provided by the Research and Technology Department of Ministry of Sciences, Tehran, Iran. For yeast culture, a medium containing (g/L) 30 g glucose, 1 g yeast extract, 0.2 K_2HPO_4 , 1 g KH_2PO_4 , 2.5 g MgSO_4 , and 9 g $(\text{NH}_4)_2\text{SO}_4$ was prepared and autoclaved for 20 min at 121°C . Once cooled, the medium was inoculated with *S. cerevisiae* followed by incubation (30°C) with shaking at 50 rpm for 20 h. The culture media was stored at 4°C for future use. The activated yeast cells were counted by the dilution method; 1 mL of seed culture was diluted in a ratio of 1/10 with sodium chloride solution, and 10 serial dilutions were obtained. Then 1 mL of each dilution was added to the solid nutrient agar medium by pour plate method [29]. Following the incubation of the plates at 30°C for 72 h, the yeast colonies were counted and showed the mean 30×10^8 CFU/mL. The yeast culture was activated daily for biosorption experiments.

Reagents. All chemicals used in this project were prepared from Merck (Darmstadt, Germany). Cd standard solution was used as the stock solution (Accu Trace, New Haven, USA). All containers were acid-washed by 15% (v/v) HNO_3 overnight, and then autoclaved at 121°C , 1.4 atm pressure, for 20 min.

Bioremoval condition. Amounts of 50 mL of milk with different metal Cd concentrations (20 , 40 , 60 , 80 , 100 mg/L)

were prepared. The effects of six variables including exposure time (1-4 days), temperature (4 and 40°C), *S. cerevisiae* concentrations (10×10^8 to 40×10^8 CFU/mL), the viability (living and non-living), shaking rate (5-20 rpm) and initial Cd concentration (20 , 40 , 60 , 80 , 100 mg/L) on the biosorption efficiency of Cd by *S. cerevisiae* were evaluated [30, 31].

Measurement of Cadmium by ICP- MS. The inductively coupled plasma mass spectrometer (ICP-MS) was used for measuring the Cd amount. First, 5 ml of milk samples were digested by using the microwave procedure at 1200 W for 10 min and then were injected to the ICP- MS spectrometer [32].

The sample mixtures containing Cd and *S. cerevisiae* were centrifuged at $2000 \times g$ for 15 min, and the supernatant recovered. Finally, Cd concentration in the supernatant was measured by using the ICP-MS. All the trials were repeated three times.

The Cd removal by *S. cerevisiae* was calculated by the following equation [33]:

$$R = \frac{C_0 - C_e}{C_0} \times 100$$

Where R is % Cd removal, C_0 is the initial concentration of Cd in milk in (mg/L), and C_e is the concentration of Cd in milk after biosorption (mg/L).

RESULTS

The effect of six variables was analyzed by the analysis of variance (ANOVA). Our analysis revealed that the Cd bioremoval was significantly affected by three variables; Cd concentration, *S. cerevisiae* concentration, and exposure time. *P*-values of less than 0.05 represent the significant variables in this project (Table 1).

Table 1. ANOVA results for the effect of parameters in Cd biosorption by *S. cerevisiae*.

Source	Sum of Squares	df	Mean Square	F-value	P-value
Shaking rate	0.2057	1	0.2057	1.92	0.3995
<i>S. cerevisiae</i> viability	0.3426	1	0.3426	3.41	0.3034
Cd concentration	8.86	1	8.86	89.62	0.0365
<i>S. cerevisiae</i> concentration	8.35	1	8.35	76.93	0.0698
Exposure time	8.76	1	8.76	81.40	0.0665
Temperature	0.3858	1	0.3858	3.23	0.3025

Effect of *S. cerevisiae* concentration on Cd bioremoval. The Cd biosorption efficiency enhanced by increasing *S. cerevisiae* concentrations from 10×10^8 to 30×10^8 CFU/mL (Fig. 1A). The maximum Cd removal (70%) was achieved at the 30×10^8 CFU/mL. By increasing the *S. cerevisiae* concentrations from 30×10^8 to 40×10^8 CFU/mL, the removal efficiency decreased.

Effect of exposure time on Cd bioremoval. During the four days of the experiment, the Cd removal by *S. cerevisiae*

increased with the extended exposure time, but it showed a faster rate on the second day and a slower trend from the second to the fourth day (Fig. 1B).

Effect of Cd concentration on the Cd bioremoval. The Cd biosorption efficiency was evaluated in different Cd concentrations (20 , 40 , 60 , 80 , and 100 mg/L). The bioremoval increased with Cd concentration from 20 to 100 mg/L. The highest biosorption (70%) was observed at the 100 mg/L of Cd (Fig. 1C).




Fig. 1. The effect of *S. cerevisiae* concentration (A), Exposure time (B) and Cd concentration (C) on Cd bioremoval

DISCUSSION

Using microorganisms as environmentally friendly biosorbents for toxic metals bioremediation from food and water resources is a novel method. In the food industry, *S. cerevisiae* is one of the potential cost-effective biosorbents for the decontamination of heavy metals from foodstuff. The biosorption occurs through a surface binding process that relates to the functional groups like hydroxyl, carboxyl, and amide groups of *S. cerevisiae* [34, 35]. In this study, we used *S. cerevisiae* for Cd removal from milk, and the effects of variables, including exposure time, *S. cerevisiae* concentration, and initial Cd concentration on the Cd bioremoval by *S. cerevisiae*, were evaluated.

The biosorption increased with *S. cerevisiae* concentration in milk samples as the active sites on the cell wall of *S. cerevisiae* became more available for attaching to the metal ions. Our results showed agreement with those of similar studies that used *S. cerevisiae* for bioremoval of heavy metals like Cd, Zinc (Zn), and Mercury (Hg) [24, 36, 37].

The Cd uptake enhanced by the increase in the exposure time from one to four days, which led to the attachment of

more Cd ions to *S. cerevisiae* receptor sites resulting in more Cd removal [38]. Our findings are in agreement with the study of Hadiani *et al.* (2018) that reported enhanced mercury bioremoval by *S. cerevisiae* when exposure time extended from 24 h to 48 h [24]. Also, Massoud *et al.* (2019) and Hatamifard *et al.* (2016) reported an increase in Pb and Hg biosorption over extended exposure time [25, 39].

The present study also assessed the bioremoval in different initial Cd concentrations, 20, 40, 60, 80, 100 mg/L with constant *S. cerevisiae* concentration (30×10^8 CFU/mL), and over the day exposure time. Clearly, by increasing the number of metal ions, their adsorption to *S. cerevisiae* receptors would increase as a result of the higher driving force for metal ions to interact with *S. cerevisiae* membrane binding sites [35, 40, 41].

The maximum Cd bioremoval (70%) with 30×10^8 CFU/mL of *S. cerevisiae* and 80 mg/L Cd concentration in milk was achieved on the 4th day. Our study provided further evidence for the suitability of *S. cerevisiae* for bioremediation of heavy metals in food industry, opening a new window for providing safe and healthy food supply.

ACKNOWLEDGMENT

We thank the National Nutrition and Food Technology Research Institute (NNFTRI) of Iran for financial support of this research project.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest associated with this manuscript.

ETHICAL APPROVAL

This article contains no assay on humans or animals by the contributing authors.

REFERENCES

1. Rajaganapathy V, Xavier F, Sreekumar D, Mandal PK. Heavy metal contamination in soil, water and Fodder and their presence in livestock and products: a review. *J Environ Sci Technol*. 2011; 4: 234-249.
2. Sardar A, Hameed K, Afzal S. Heavy metals contamination and what are the impacts? *Greener J Environ Manag Public Saf*. 2013; 2: 17-19.
3. Patton S. Milk Its remarkable contribution to human health and well-being. Taylor and Francis Publishers; 2017; 250-288.
4. Maslehat S, Mostafavi E. The History of Milk. *J Med Microbiol Infect Dis*. 2018; 6 (4): 87-90.
5. Weldelessi T, Balwant H, Oves M. Chemical Contaminants in Soil. Air and Aquatic Ecosystem Mod Age *Environ Prob Rem*. 2017; 25: 1-22.
6. Awasthi V, Bahman S, Thakur LK, Singh SK, Dua A, Ganguly S. Contaminants in milk an assessment study. *Indian J Public Health*. 2012; 56: 25-40.
7. Lokhande RS, Singare PU, Pimple DS. Toxicity Study of Heavy Metals Pollutants in Waste Water Effluent Samples Collected from Talaja Industrial Estate of Mumbai, India. *Res Environ*. 2011; 1 (1): 13-19.
8. Li XG, Li DB, Yan ZN, Ao YS. Biosorption and bioaccumulation characteristics of cadmium by plant growth-promoting rhizobacteria. *RSC Adv*. 2018; 83-92.
9. Sari A, Tuzen M. Biosorption of Pb (II) and Cd (II) from aqueous solution using green alga (*Ulva lactuca*) biomass. *J Hazard Mater*. 2008; 152: 302-308.
10. Hajjaligol S, Taher MA, Malekpour A. A new method for the selective removal of cadmium and zinc ions from aqueous solution by modified clinoptilolite. *Adsorp Sci Technol*. 2006; 24: 487-496.
11. Alani MS, Al-Azzawi MN. Assessment of Lead Cadmium and Copper concentrations in Raw Milk Collected from different location in Iraq. *Iraq J Sci*. 2016; 56 (1): 350-355.
12. Ayar A, Sert D, Akin N. The trace metal levels in milk and dairy products consumed in middle Anatolia-Turkey. *Environ Monitor Assess*. 2015; 152 (1):1-12.
13. Starska k, Wojciechowska-Mazurek M, Mania M, Brulinska-Ostrowska E, Biernat U, Karlowski K. Noxious elements in milk and milk products in Poland. *Polish J Environ*. 2011; 20 (4): 1043-1051.
14. Qin L, Wang X, Li W, Tong X, Tong W. The minerals and heavy metals in cow's milk in China *Japan J Health Sci*. 2009; 55 (2): 300-305.
15. Nejatolahi M, Mehrjo F, Sheykhi A, Bineshpour M. Lead Concentrations in Raw Cows' Milk from Fars Province of Iran. *Americ J Food Nut*. 2014; 2 (5): 92-94.
16. Najarneshad V, Akbarabadi M. Heavy metals in raw cow and ewe milk from north-east Iran. *Food Add Contamin*. 2013; 12: 2-6.
17. Xu J, Bravo AJ, Lagerkvist A, Bertilsson A, Sjöblom R, Kumpiene J. Sources and remediation techniques for metal contaminates. *Environ Int*. 2015; 74: 42-53.
18. Pakdel M, Soleimani-Zad S, Akbari-Alavijeh S. Screening of Lactic acid bacteria to detect potent biosorbents of lead and cadmium. *Food Control*. 2018; 28: 33-40.
19. Kumar N, Kumar V, Panwar R, Ram C. Efficacy of indigenous probiotic *Lactobacillus* strains to reduce cadmium bioaccessibility - An in vitro digestion model. *Environ Sci Pollut Res*. 2016; 25: 22-33.
20. Hadiani MR, Khosravi-Darani K, Rahimifard N, Younesi H. Biosorption of low concentration levels of Lead (II) and Cadmium (II) from aqueous solution by *Saccharomyces cerevisiae*. *Biocat Agr Biotech*. 2018; 15: 25-34.
21. Massoud R, Hadiani MR, Khosravi Darani K. Bioremediation of heavy metals in food industry Application of *Saccharomyces cerevisiae*. *Electron J Biotechnol*. 2019; 37: 56-60.
22. Salimi M, Mahzounieh M. *Saccharomyces cerevisiae* on Mo and Cd removal. *J Med Microbiol Infect Dis*. 2015; 3 (2): 18-22.
23. Amirnia S, Ray MB, Margaritis A. Heavy metals removal from aqueous solutions using *Saccharomyces cerevisiae* in a novel continuous bioreactor-biosorption system. *Chem Eng J*. 2015; 264: 863-872.
24. Hadiani MR, Khosravi-Darani K, Rahimifard N, Younesi H. Assessment of Mercury biosorption by *Saccharomyces cerevisiae* Response surface methodology for optimization of low Hg (II) concentrations. *J Environ Chem Eng*. 2018; 6: 4980-4987.
25. Massoud R, Khosravi-Darani K, Sharifan A, Asadi GH. Lead Bioremoval from Milk by *Saccharomyces cerevisiae*. *Biocat Agri Biotech*. 2019; 22: 11-20.
26. Infante C, Arco D, Angulo E. Removal of lead mercury and nickel using the yeast *Saccharomyces cerevisiae*. *Rev MVZ Cordoba*. 2014; 19: 4141-4149.
27. Chen C, Wang J. Review on biosorption of heavy metal by *Saccharomyces cerevisiae*. *Chin J Biotechnol*. 2006; 26: 69-76.
28. Farhan SN, Khadom AA. Biosorption of heavy metals from aqueous solutions by *Saccharomyces Cerevisiae*. *Int J Ind Chem*. 2015; 6: 119-130.
29. Iranian National Standardization Organization, Microbiology of milk and milk products – Specifications and test methods. *ISO 2406*: 2017.
30. Sepehr M.N. Zarrabi M. Amrane A. Removal of CR (III) from model solutions by isolated *Aspergillus Niger* and *Aspergillus*

oryzae living microorganisms: Equilibrium and kinetic studies. J Taiwan Inst Chem Eng. 2012; 43: 420-427.

31. Wang T, Sun H, Jiang C, Mao H, Zhang Y. Immobilization of Cd in soil and changes of soil microbial community by bioaugmentation of UV-mutated *Bacillus subtilis* 38 assisted by biostimulation. Eur J Soil Biol. 2014; 65: 62-69.

32. Khan N, Jeong S, Hwang M, Kim JS, Choi SH, Yeong E, et al. Analysis of minor and trace elements in milk and yogurts by inductively coupled plasma-mass spectrometry (ICP-MS). Food Chem. 2014; 147: 220-224.

33. Goksungur Y, Uren S, Guvenc U. Biosorption of cadmium and lead ions by ethanol treated waste baker's yeast biomass. Bioresour Technol. 2005; 96: 103-109.

34. Wang JL, Chen C. Biosorption of heavy metals by *Saccharomyces cerevisiae* a review. Biotechnol Adv. 2006; 24: 427-451.

35. Zoghi A, Khosravi-Darani K, Sohrabvandi S. Surface Binding of Toxins and Heavy Metals by Probiotics. Mini Rev Med Chem. 2014; 14: 84-98.

36. Vasudevan P, Padmavathy V, Dhingra SC. Kinetics of biosorption of cadmium on Baker's yeast. Bioresour Technol. 2003; 89: 281-287.

37. Vinopal S, Ruml T, Kotrba P. Biosorption of Cd and Zn by cell surface-engineered *Saccharomyces cerevisiae*. Int Biodeterior Biodegrad. 2007; 60: 96-102.

38. Amini A, Younesi H. Biosorption of Cd (II) Ni (II) and Pb (II) from Aqueous Solution by Dried Biomass of *Aspergillus niger*, Application of Response Surface Methodology to the Optimization of Process Parameters. Clean Soil Air Water J. 2009; 37: 776-786.

39. Hatami Fard G, Mehrnia MR. Investigation of mercury removal by Micro-Algae dynamic membrane bioreactor from simulated waste water. J Environ Chem Eng. 2016; 10: 25-33.

40. Saber-Samandari S, Gazi M. Removal of mercury (II) from aqueous solution using chitosan-graft-Polyacrylamide Semi-IPN hydrogels. Sep Sci Technol. 2013; 48: 1382-1390.

41. Kazy SKD, Souza SF, Sar P. Uranium and thorium sequestration by a *Pseudomonas* sp. mechanism and chemical characterization. J Hazard Mater. 2009; 163: 65-72.

Cite this article:

Massoud R, Sharifan A, Khosravi-Darani K, Asadi GH. Cadmium Bioremoval by *Saccharomyces cerevisiae* in Milk. J Med Microbiol Infect Dis, 2020; 8 (1): 29-33. DOI: 10.29252/JoMMID.8.1.29.